

Marginal Deformations of $\mathcal{N} = 4$ SYM and Open vs. Closed String Parameters

Manuela Kulaxizi

*C. N. Yang Institute for Theoretical Physics
State University of New York at Stony Brook
Stony Brook, NY 11794-3840, USA*

E-mail: kulaxizi@insti.physics.sunysb.edu

ABSTRACT

We construct the gravity dual to the generic Leigh–Strassler deformation of $\mathcal{N} = 4$ SYM as well as the corresponding deformed flat space geometry up to third order in the deformation parameter ρ . We achieve this by first determining the set of open string parameters $(\mathcal{G}, \Theta)$ encoding the deformation of the moduli space and then mapping them to the closed string ones (g, B) . Our method is thus almost purely algebraic involving the supergravity equations of motion only as a consistency check. We trace the main reason for the discrepancy at higher orders in the deformation parameter to the nonassociative nature of the noncommutativity matrix Θ .

1 Introduction

The AdS/CFT correspondence [1, 2, 3] offers an equivalence between gauge theory and gravity. In its original form, relates superconformal $\mathcal{N} = 4$ SU(N) Super Yang–Mills to closed string theory on $\text{AdS}_5 \times \text{S}^5$ with N units of RR–flux. While closed strings on nontrivial backgrounds with RR–fluxes are still in many ways intractable, their low energy description in terms of supergravity is not. From the gauge theory point of view, this limit corresponds to large N and strong t’Hooft coupling λ . This makes the correspondence extremely useful in that it provides a window into understanding the physics of gauge theories in a region that is otherwise difficult to explore. By now the original proposal has been greatly extended so as to cover gauge theories with less amount of supersymmetry and/or a running coupling constant [4, 5, 6, 7, 8, 9]. Despite however the progress made thus far the natural question of how to specify the gravitational background corresponding to a given gauge theory, is still far from being understood ¹. In this article we will attempt to give an alternative point of view on this matter while investigating marginal deformations of the $\mathcal{N} = 4$ theory.

Supersymmetric deformations of the original AdS/CFT proposal have been extensively explored with the main cases of interest being, exactly marginal and relevant deformations. Whereas it would have been natural to think that the gravity dual backgrounds of the former would be more accessible than those of the latter it actually turned out the opposite. In fact the gravity duals of a class of supersymmetric mass deformations, all along their renormalization group flow, were discovered quite early on — see for example [16, 17, 18] and references therein — even though in most cases an analytic solution was provided only for the associated conformal fixed points. The main reason is that these backgrounds can be analyzed using the truncation to five–dimensional supergravity, something which is not possible for marginal deformations of the $\mathcal{N} = 4$ theory. Actually it was only fairly recently that the authors of [19] succeeded in constructing the corresponding backgrounds for a subclass of these latter theories.

Marginal deformations of $\mathcal{N} = 4$ SYM preserve $\mathcal{N} = 1$ supersymmetry and are mainly described by two parameters, denoted as β and ρ , in addition to the gauge coupling g_{YM} . In [19] Lunin and Maldacena discovered the geometry dual to the β –deformed theory i.e. when $\rho = 0$. In this case, extra U(1) global symmetries are present which played a significant part in the construction of the new solution. When $\rho \neq 0$ however, the theory does not preserve any continuous symmetries other than the U(1) R–symmetry and the problem has resisted solution thus far. In this note, we construct the supergravity solution dual to the ρ –deformed theory up to third order in the deformation parameter. We propose a method which — although rudimentary at this stage — may provide additional means in exploring deformations of the original AdS/CFT proposal. The main idea, is to encode information from the gauge theory moduli space into the set of *open* string parameters $(\mathcal{G}, \Theta)$ and then map them to the *closed* string ones (g, B) .

In [20] we achieved the first step in this direction by establishing a precise relation between transverse space noncommutativity and β –deformations of $\mathcal{N} = 4$ SYM. We identified the role of the noncommutativity matrix Θ in the solution generating technique of Lunin and Maldacena and showed how to explicitly construct it, relying on gauge theory data and basic notions of AdS/CFT. The resulting noncommutativity matrix Θ , along with the flat metric of the transverse space $\mathcal{G}_{flat}$, provided an alternative way of encoding the moduli space. They constitute the *open* string parameters $(\mathcal{G}, \Theta)$. Following here the method of [20] we acquire the corresponding set of open string data for the ρ –deformed gauge theory. In this case, it turns out that Θ defines a *nonassociative*, as well as noncommutative, deformation of the transverse space.

Having obtained the open string parameters, we move on to explore possible mappings to the closed string ones. It is natural to expect by T–duality, that the results of Seiberg and Witten [21] — see also [22] — relating noncommutativity to the presence of a nonvanishing B–field, will be valid independently of whether the noncommutative deformation is along or transverse to the D–brane. It follows that the equations which establish a connection between the open $(\mathcal{G}, \Theta)$ and closed (g, B) string fields in that setting, present a potential mapping of the associated parameters in this case as well. We therefore employ them in the context of the ρ –deformation and determine the deformed flat space background into where D–branes are immersed up to third order in the deformation parameter.

¹For an existing approach to related issues see for instance [10, 11, 12, 13, 14, 15] and references therein.

We then consider the effective action of the ρ -deformed gauge theory, obtained by giving a vacuum expectation value to one of the scalars and integrating out the massive fields. According to [23, 24, 25, 26, 27, 28], the leading IR large N part of this action should coincide with the DBI action for a D3-brane immersed in the dual background. We observe that in the case of the β -deformed gauge theory, the corresponding DBI action is characterized by the open string data $(\mathcal{G}_{AdS_5 \times S^5}, \Theta)$ and that the associated NS-NS closed string fields (g, B) are part of the *exact* Lunin-Maldacena solution. This is not surprising. Indeed, the Lagrangian description of this theory can be given in terms of the $\mathcal{N} = 4$ Lagrangian with the product of matter fields replaced by a star product of the Moyal type. Subsequently, all amplitudes in the planar limit can be shown [29] to be proportional up to a phase to their $\mathcal{N} = 4$ counterparts. Then the open string data $(\mathcal{G}_{AdS_5 \times S^5}, \Theta = 0)$ of the $\mathcal{N} = 4$ SYM theory are naturally promoted to the set $(\mathcal{G}_{AdS_5 \times S^5}, \Theta)$. Can something similar occur for the ρ -deformation?

Since the noncommutativity parameter now defines a nonassociative deformation, it is hard to imagine how the planar equivalence with the parent $\mathcal{N} = 4$ theory could be achieved. Yet nonassociativity is a second-order in ρ effect and it turns out that up to third order in ρ this is indeed the case. In other words, mapping the open string fields to the closed ones provides again a supergravity solution to this order — the gravity dual of the ρ -deformed gauge theory.

The structure of this paper is the following: In section 2 we review some known facts about marginal deformations of the $\mathcal{N} = 4$ theory and their gravity duals. In addition, we explore some special points in the deformation space for which the general theory with $\beta \neq 0$ and $\rho \neq 0$ is equivalent to an exactly marginal deformation with either $\tilde{\beta} = 0$ or $\tilde{\rho} = 0$. In section 3, we employ the method proposed in [20] in order to determine the relevant noncommutativity matrix for the ρ -deformation. Having acquired the open string data, we move on to map them to the closed string fields (g, B) . This procedure is illustrated in section 4 where we derive the ρ -deformed flat space geometry up to third order in the deformation parameter. In section 5 we proceed with considerations on the DBI action which provide us with the gravity dual of the ρ -deformed theory to the same order. Section 6, contains an extensive discussion on the ideas set forth in this note, whereas conclusions and possible extensions are given in section 7.

2 The Leigh Strassler Deformation

Not long after it was realized that $\mathcal{N} = 4$ SU(N) Super Yang Mills theory is finite (see e.g. [30] for an account), it became clear that it might not be the only four dimensional theory with that property [31, 32, 33, 34, 35]. It was however almost ten years later, when Leigh and Strassler undertook a systematic study of marginal deformations of $\mathcal{N} = 4$ and indeed showed that there exists a whole class of $\mathcal{N} = 1$ supersymmetric gauge theories satisfying both the requirements for conformal invariance and finiteness [36]. More precisely, they showed that the $\mathcal{N} = 4$ theory admits a three-complex-parameter family of marginal deformations which preserve $\mathcal{N} = 1$ supersymmetry and are described by the following superpotential:

$$\mathcal{W} = ih \text{Tr} \left[(e^{i\beta} \Phi_1 \Phi_2 \Phi_3 - e^{-i\beta} \Phi_1 \Phi_3 \Phi_2) + \rho (\Phi_1^3 + \Phi_2^3 + \Phi_3^3) \right] \quad (1)$$

where Φ^I with $I = 1, 2, 3$ are the three chiral superfields of the theory. Together with the gauge coupling g_{YM} , the complex parameters (h, β, ρ) that appear in the superpotential constitute the four couplings of the theory.

While it is clear at the classical level that these deformations are marginal — since all operators of the component Lagrangian have classical mass dimension equal to four — this is not necessarily true quantum mechanically. Leigh and Strassler realized that by using the constraints of $\mathcal{N} = 1$ supersymmetry and the exact NSVZ beta-functions [37, 38, 39] written in terms of the various anomalous dimensions of the theory, it was possible to express the conditions for conformal invariance of the quantum theory, through linearly dependent equations which were therefore likely to have nontrivial solutions. In this way, they were able to demonstrate that the deformation of (1) is truly marginal at the quantum level, so long as the four couplings of the theory satisfy a single complex constraint $\gamma(g_{YM}, \kappa, \beta, \rho) = 0$. In other words, there exists a three-complex-dimensional surface $\gamma(g_{YM}, \kappa, \beta, \rho) = 0$ in the space of couplings, where both beta functions and anomalous dimensions vanish and thus the $\mathcal{N} = 1$ gauge theories mentioned above are indeed conformally

invariant. In general, the function γ is not known beyond two-loops [40, 41, 42, 43, 44] in perturbation theory, where it reads:

$$|h|^2 \left[\frac{1}{2} \left(|q|^2 + \frac{1}{|q|^2} \right) - \frac{1}{N^2} \left| q - \frac{1}{q} \right|^2 + |\rho|^2 \left(\frac{N^2 - 4}{2N^2} \right) \right] = g_{YM}^2 \quad (2)$$

with q defined as $q = e^{i\beta}$ and N the number of colours of the gauge theory.

For the β -deformed gauge theory, *i.e.*, obtained by setting $\rho = 0$ in the superpotential of equation (1), the Leigh–Strassler constraint at two loops can be written as:

$$|h|^2 \left[\frac{1}{2} \left(|q|^2 + \frac{1}{|q|^2} \right) - \frac{1}{N^2} \left| q - \frac{1}{q} \right|^2 \right] = g_{YM}^2 \quad (3)$$

In this case, one immediately notices that when $\beta = \beta_{\mathbb{R}} \in \mathbb{R}$ therefore $|q| = 1$, (3) reduces to:

$$|h|^2 \left[1 - \frac{1}{N^2} \left| q - \frac{1}{q} \right|^2 \right] = |h|^2 \left(1 - \frac{4}{N^2} \sin^2 \beta_{\mathbb{R}} \right) = g_{YM}^2 \quad (4)$$

which in the large N limit yields: $|h|^2 = g_{YM}^2$. Despite the fact that this result was obtained from the two-loop expression of the conformal invariance condition, it has been shown to be true to all orders in perturbation theory at the planar limit [29] (see also [45, 44]). Actually the author of [29] went even further and showed that all planar amplitudes in the $\beta = \beta_{\mathbb{R}} \in \mathbb{R}$ theory are proportional to their $\mathcal{N} = 4$ counterparts, thus explicitly proving finiteness and conformal invariance. The proof made use of an existing proposal [19] for an equivalent "noncommutative" realization of the theory. For the more general case of complex $\beta = \beta_{\mathbb{R}} + i\beta_{\mathbb{I}}$, equation (3) in the planar limit reads:

$$\frac{1}{2} |h|^2 \left(|q|^2 + \frac{1}{|q|^2} \right) = |h|^2 \cosh(2\beta_{\mathbb{I}}) = g_{YM}^2 \quad (5)$$

It is then evident that the coupling constant h receives corrections with respect to its $\mathcal{N} = 4$ SYM value. Nevertheless, diagrammatic analysis [29] showed that all planar amplitudes with external gluons are equal to those of the $\mathcal{N} = 4$ theory up to a five-loop level. To this order and beyond, it is most likely that the planar equivalence between the parent theory and its deformation will break down. For (more) recent investigation on β -deformations from the gauge theory point of view see [43, 44, 46, 47, 48, 49, 50].

Special points along the deformation occur when β is a root of unity. These points have been studied early on [51, 52] with a dual interpretation as orbifolds with discrete torsion. The marginally deformed theories have been further explored in [53, 54, 55], and several remarkable properties have been demonstrated. In particular it was shown that as expected, the S-duality of $\mathcal{N} = 4$ extends to their space of vacua, and that, again for special values of β , there are also new Higgs branches on moduli space. These are mapped by S-duality to completely new, confining branches which appear only at the quantum level. Furthermore, at large N the Higgs and confining branches can be argued to be described by Little String Theory [55]. Finally, the possibility of an underlying integrable structure for the deformed theories in analogy with $\mathcal{N} = 4$ SYM, was investigated at special values of the deformation parameter in [56, 57] and for generic β in [58, 59, 60, 61].

2.1 Marginal deformations and gauge/gravity duality

A natural place to explore theories that arise as marginal deformations of $\mathcal{N} = 4$ SU(N) SYM is the AdS/CFT correspondence where the strong coupling regime of the undeformed theory is realized as weakly coupled supergravity on $\text{AdS}_5 \times S^5$. Due to superconformal symmetry, the dual gravitational description of these theories is expected to be of the form: $\text{AdS}_5 \times \tilde{S}^5$ with $\tilde{S}^5$ a sphere deformed by the presence of additional NS–NS and RR fluxes. Indeed in [62], where the dual background was constructed to second order in the deformation parameters, it was shown that apart from the already present five-form flux one should also turn on (complexified) three-form flux $G_{(3)}$ along the S^5 .

Essential progress however in this direction was only recently achieved through the work of Lunin and Maldacena [19]. The authors of [19] succeeded in finding the *exact* gravity dual of the β -deformed gauge theory.

In this case, apart from the $U(1)_R$ R-symmetry the theory preserves two global $U(1)$ s, which act on the superfields in the following way:

$$\begin{aligned} U(1)_1 : \quad & (\Phi_1, \Phi_2, \Phi_3) \rightarrow (\Phi_1, e^{i\alpha_1} \Phi_2, e^{-i\alpha_1} \Phi_3) \\ U(1)_2 : \quad & (\Phi_1, \Phi_2, \Phi_3) \rightarrow (e^{-i\alpha_2} \Phi_1, e^{i\alpha_2} \Phi_2, \Phi_3) \end{aligned} \quad (6)$$

The main idea underlying the solution generating technique proposed in [19], was the natural expectation that the two $U(1)$ symmetries preserved by the deformation would be realized geometrically in the dual gravity solution. For $\beta = \beta_{\mathbb{R}} \in \mathbb{R}$ their prescription amounts to performing an $SL(2, \mathbb{R})$ transformation on the complexified Kähler modulus τ of the two torus associated with the $U(1)$ symmetries in question. The specific element of $SL(2, \mathbb{R})$ under consideration is: $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \equiv \begin{pmatrix} 1 & 0 \\ c & 1 \end{pmatrix}$. It is chosen so as to ensure that the new solution will present no singularities as long as the original one is non-singular and its sole free parameter c is naturally identified with the real deformation parameter $\beta_{\mathbb{R}}$ of the gauge theory.

Later on, the method of Lunin and Maldacena was reformulated [63] in terms of the action of a T-duality group element on the background matrix $E = g + B$ providing a significantly easier way of obtaining the new solutions. In particular, it was shown [63] that one can embed the $SL(2, \mathbb{R})$ that acts on the Kähler modulus into the T-duality group $O(3, 3, \mathbb{R})$ in the following way:

$$\mathcal{T} = \begin{pmatrix} \mathbf{1} & \mathbf{0} \\ \mathbf{\Gamma} & \mathbf{1} \end{pmatrix} \quad \text{where now} \quad \mathbf{\Gamma} \equiv \begin{pmatrix} 0 & -\beta_{\mathbb{R}} & \beta_{\mathbb{R}} \\ \beta_{\mathbb{R}} & 0 & -\beta_{\mathbb{R}} \\ -\beta_{\mathbb{R}} & \beta_{\mathbb{R}} & 0 \end{pmatrix} \quad (7)$$

where $\mathbf{1}$ and $\mathbf{0}$ represent the 3×3 identity and zero matrices respectively. Suppose then that $E_0 = g_0 + B_0$ denotes the part of the original supergravity background along the $U(1)$ isometry directions which are to be deformed. Acting on E_0 with the T-duality group element $\mathcal{T}$ of (7) one obtains the NS-NS fields of the deformed solution in terms of E_0 and $\mathbf{\Gamma}$ according to:

$$\begin{aligned} E &= \frac{1}{E_0^{-1} + \mathbf{\Gamma}} \\ e^{2\Phi} &= e^{2\Phi_0} \det(1 + E_0 \mathbf{\Gamma}) \equiv e^{2\Phi_0} G \end{aligned} \quad (8)$$

The RR-fields of the background can be computed using the T-duality transformation rules of [64, 65, 66, 67, 68], however the details of this transformation need not concern us here. As an example, let us consider ten-dimensional flat space parametrized as:

$$ds^2 = -dt^2 + \sum_{\mu=1}^3 dx^\mu dx_\mu + \sum_{i=1}^3 (dr_i^2 + r_i^2 d\varphi_i^2) \quad (9)$$

In this case E_0 will contain the components of the flat metric along the polar angles φ_i . Applying equations (8) yields:

$$\begin{aligned} ds^2 &= -dt^2 + \sum_{\mu=1}^3 dx^\mu dx_\mu + \sum_{i=1}^3 (dr_i^2 + G r_i^2 d\varphi_i^2) + \beta_{\mathbb{R}} G r_1^2 r_2^2 r_3^2 \left(\sum_{i=1}^3 d\varphi_i \right)^2 \\ e^{2\Phi} &= G, \quad G^{-1} = 1 + \beta_{\mathbb{R}}^2 \left(\sum_{i \neq j} r_i^2 r_j^2 \right), \quad B = \beta_{\mathbb{R}} G \left(\sum_{i \neq j} r_i^2 r_j^2 d\varphi_i d\varphi_j \right) \end{aligned} \quad (10)$$

This is the deformed flat space geometry where by placing D3-branes at the origin and taking the near horizon limit, one obtains the gravity dual to the β -deformed gauge theory. Alternatively, the latter background can

be constructed by applying (8) on $\text{AdS}_5 \times \text{S}^5$ representing the dual gravitational description of the undeformed parent $\mathcal{N} = 4$ theory:

$$\begin{aligned}
ds^2 &= R^2(ds_{\text{AdS}_5}^2 + ds_5^2), \quad \text{where :} \quad ds_5^2 = \sum_i (d\mu_i^2 + G\mu_i^2 d\varphi_i^2) + \hat{\beta} G \mu_1^2 \mu_2^2 \mu_3^2 \left(\sum_i d\varphi_i \right)^2 \\
e^{2\Phi} &= e^{2\Phi_0} G, \quad G^{-1} = 1 + \hat{\beta}^2 \left(\sum_{i \neq j} \mu_i^2 \mu_j^2 \right), \quad \hat{\beta} = R^2 \beta_{\mathbb{R}}, \quad R^4 = 4\pi e^{\Phi_0} N \\
B &= \hat{\beta} R^2 G \left(\sum_{i \neq j} \mu_i^2 \mu_j^2 d\varphi_i d\varphi_j \right) \quad C_2 = -\beta_{\mathbb{R}} (16\pi N) \omega_1 \left(\sum_i d\varphi_i \right) \\
F_5 &= (16\pi N) (\omega_{\text{AdS}_5} + G\omega_{\text{S}^5}), \quad \omega_{\text{S}^5} = d\omega_1 d\varphi_1 d\varphi_2 d\varphi_3, \quad \omega_{\text{AdS}_5} = d\omega_4
\end{aligned} \tag{11}$$

Reformulating the Lunin–Maldacena generating solution technique in terms of the T–duality group action, made especially transparent its relation to similar methods employed in the context of noncommutative gauge theories². In [20] we identified the prescriptions used in these cases and particularly showed that matrix Γ of (7) is precisely the noncommutativity matrix Θ associated with the deformation of the transverse space. Moreover, we proposed a method for determining Θ based solely on information from the gauge theory Lagrangian and basic notions of AdS/CFT. The main idea was to think of the matter fields as coordinates $(z^I, \bar{z}^{\bar{I}})$ parametrizing the space transverse to the D3–brane where the gauge theory lives. Then, reality properties, global symmetries and marginality constrained the form of Θ to be:

$$\Theta_{\beta} = a \begin{pmatrix} 0 & z_1 z_2 & -z_1 z_3 & 0 & -z_1 \bar{z}_2 & z_1 \bar{z}_3 \\ -z_1 z_2 & 0 & z_2 z_3 & \bar{z}_1 z_2 & 0 & -z_2 \bar{z}_3 \\ z_3 z_1 & -z_2 z_3 & 0 & -\bar{z}_1 z_3 & \bar{z}_2 z_3 & 0 \\ 0 & -\bar{z}_1 z_2 & \bar{z}_1 z_3 & 0 & \bar{z}_1 \bar{z}_2 & -\bar{z}_1 \bar{z}_3 \\ z_1 \bar{z}_2 & 0 & -\bar{z}_2 z_3 & -\bar{z}_1 \bar{z}_2 & 0 & \bar{z}_2 \bar{z}_3 \\ -\bar{z}_3 z_1 & \bar{z}_3 z_2 & 0 & \bar{z}_1 \bar{z}_3 & -\bar{z}_2 \bar{z}_3 & 0 \end{pmatrix} \tag{12}$$

with $a = 2 \sin \beta_{\mathbb{R}}$. While it may seem that Θ_{β} is dissimilar from matrix Γ of (7), a coordinate transformation from $(z^I, \bar{z}^{\bar{I}})$ to polar coordinates (r_i, φ_i) on $\mathbb{R}^6$ reveals that:

$$\Theta_{\beta} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -a & a \\ 0 & 0 & 0 & a & 0 & -a \\ 0 & 0 & 0 & -a & a & 0 \end{pmatrix} \tag{13}$$

thereby proving their identification³.

For the general case of complex β , to obtain the dual background, one needs to perform an additional $SL(2, \mathbb{R})_s$ transformation on the solution corresponding to $\beta_{\mathbb{R}}$. By $SL(2, \mathbb{R})_s$ we denote here the $SL(2, \mathbb{R})$ symmetry of ten dimensional type IIB supergravity which acts nontrivially on the complexified scalar and two–form fields of the theory. Being a symmetry of the equations of motion it can be used to generate distinct solutions. Subsequent work on the subject of the β –deformed gauge theories has provided further checks of the AdS/CFT correspondence [71, 72, 73, 74, 75, 76] whereas generalizations as well as applications of the solution generating technique introduced in [19] were considered in [77, 59, 76, 78, 79, 80].

2.2 Special points along the general Leigh–Strassler deformation

In this article we will be mainly interested in the ρ –deformed gauge theories. In this case — when $\rho \neq 0$ — the theory does not preserve additional $U(1)$ symmetries, it is however invariant under a global discrete

²In fact, evidence relating marginal deformations and noncommutativity was given earlier both at strong [52] and weak [69, 70] coupling.

³Note that the Lunin–Maldacena technique is valid for small $\beta_{\mathbb{R}}$ in which case $a = 2\beta_{\mathbb{R}}$

symmetry $\mathbb{Z}_3 \times \mathbb{Z}_3$ acting on the superfields as:

$$\begin{aligned}\mathbb{Z}_{3(1)} : \quad & (\Phi_1, \Phi_2, \Phi_3) \rightarrow (\Phi_3, \Phi_1, \Phi_2) \\ \mathbb{Z}_{3(2)} : \quad & (\Phi_1, \Phi_2, \Phi_3) \rightarrow (\Phi_1, e^{\frac{i2\pi}{3}} \Phi_2, e^{-\frac{i2\pi}{3}} \Phi_3)\end{aligned}\tag{14}$$

As previously mentioned, the presence of global $U(1)$ s is crucial in the solution generating technique of Lunin and Maldacena which is therefore not applicable here. In fact, the exact gravity dual for this case is still unknown. Despite however that the absence of extra continuous symmetries makes the cases of $\rho = 0$ and $\rho \neq 0$ radically different, there exist special points along the space of couplings where the two theories are not only similar but actually equivalent.

As first pointed out in [52] — see also [57, 81] — it is possible to start with either the set $(\beta, \rho) = (\beta, 0)$ or $(\beta, \rho) = (0, \rho)$, and via a field redefinition reach a point in the deformation space with $(\tilde{\beta} \neq 0, \tilde{\rho} \neq 0)$. The final point will obviously not represent the most general deformation, since the new couplings $\tilde{\beta}$ and $\tilde{\rho}$ will be given in terms of the original parameter. In other words, there will exist a function $f(\tilde{\beta}, \tilde{\rho}) = 0$ relating the two. Furthermore, requiring that the field redefinition be the result of a unitary transformation imposes a restriction on the original value of the coupling; be it β or ρ . In particular, suppose that we consider the marginally deformed theory at the point $(\beta, \rho = 0)$ and then take:

$$\begin{pmatrix} \Phi_1 \\ \Phi_2 \\ \Phi_3 \end{pmatrix} \rightarrow \begin{pmatrix} A & A & A \\ B & \omega B & \omega^2 B \\ C & \omega^2 C & \omega C \end{pmatrix} \begin{pmatrix} \Phi_1 \\ \Phi_2 \\ \Phi_3 \end{pmatrix}\tag{15}$$

with Φ_I the three chiral superfields and $\omega = e^{i2\pi/3}$ the third root of unity. Note here that since the deformation enters only in the superpotential, it suffices to consider transformations that affect the chiral fields independently from the antichiral ones. In other words, we do not expect mixing between holomorphic and antiholomorphic pieces. If we furthermore impose the following conditions on the free parameters A, B, C : $|A| = |B| = |C| = \frac{1}{\sqrt{3}}$ and $ABC = \pm \frac{i\lambda}{3\sqrt{1+2\cos 2\beta}}$ with $\lambda \in \mathbb{C}$, we find that the original β -deformed gauge theory is equivalent to the marginally deformed $\mathcal{N} = 4$ SYM theory with coupling constants:

$$\tilde{\rho} = \pm \frac{2 \sin \beta}{3\sqrt{1+2\cos 2\beta}} \quad \text{and} \quad e^{i\tilde{\beta}} = \pm \frac{2 \cos(\beta - \frac{\pi}{6})}{\sqrt{1+2\cos 2\beta}}\tag{16}$$

provided that $\beta = \beta_{\mathbb{R}} + i\beta_{\mathbb{I}}$ satisfies the following equation:

$$4 \cos 2\beta_{\mathbb{R}} \cos 2\beta_{\mathbb{I}} + 4 \cos^2 2\beta_{\mathbb{R}} + 4 \cos^2 2\beta_{\mathbb{I}} - 3(1 + 3\lambda) = 0\tag{17}$$

Solutions to (17) define special regions in the coupling constant space where the Leigh–Strassler theory with generic β and $\rho = 0$ is equivalent to a theory with both $\tilde{\beta}$ and $\tilde{\rho}$ nonvanishing but constrained to satisfy a specific relation dictated from (16). It is worth remarking here that there is no solution of (16) and (17) for which *both* β and $\tilde{\beta}$ are real. This is particularly interesting, because it is only for the β -deformed gauge theory with $\beta = \beta_{\mathbb{R}} \in \mathbb{R}$ that a precise connection with noncommutativity is possible. It is natural to wonder whether distinct unitary field redefinitions of a type similar to (15) could take us from different β 's to different $\tilde{\beta}$ and $\tilde{\rho}$. It is however not hard to deduce that up to a phase in $\tilde{\rho}$ — which can be reabsorbed in the definition of the coupling constant h — and a sign in $\tilde{\beta}$, all such unitary transformations share the same starting point (17) and lead to the same theory (16).

In an analogous manner one can find specific values of ρ for which the theory with $\beta = 0$ is equivalent to another one with both couplings $\tilde{\beta}$ and $\tilde{\rho}$ turned on. Detailed analysis in this case shows in fact that such a mapping is possible for *any* original value of ρ with parameters $\tilde{\rho}$ and $\tilde{\beta}$ given by:

$$\tilde{\rho}^2 = -\frac{\rho^2}{\rho^2 + 3}, \quad \text{and} \quad \sin^2 \tilde{\beta} = -\tilde{\rho}^2 = \frac{\rho^2}{\rho^2 + 3}\tag{18}$$

The precise field redefinition through which this is achieved, is of the form of (15):

$$\begin{pmatrix} \Phi_1 \\ \Phi_2 \\ \Phi_3 \end{pmatrix} \rightarrow \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix} \begin{pmatrix} \Phi_1 \\ \Phi_2 \\ \Phi_3 \end{pmatrix}\tag{19}$$

Note here again that $\tilde{\beta} = \tilde{\beta}_{\mathbb{R}} \in \mathbb{R}$ if and only if $\rho \in \mathbb{R}$ which implies that $\tilde{\rho} \in \mathbb{I}$. If one additionally assumes that $\tilde{\beta}_{\mathbb{R}} \in \mathbb{R} \ll 1$ then the deformed theory with $\beta = 0$ and $\rho = q_1 \in \mathbb{R}$ is equivalent to a theory with $2\sin\beta = \pm 2\frac{q_1}{\sqrt{3}}$ and $\tilde{\rho} = \pm i\frac{q_1}{\sqrt{3}} \in \mathbb{I}$. In section 3, we will see that this particular point in the deformation space naturally shows up in the noncommutative description of the moduli space. This will provide us with a non-trivial check on the consistency of the noncommutative interpretation.

So far we have looked at special points in the space of couplings which can be studied at the level of the gauge theory lagrangian. There are however a couple of interesting observations one can additionally make on the basis of the Leigh–Strassler constraint as this is given in equation (2). Notice first that (2) reduces in the planar limit to:

$$|h|^2 \left[\frac{1}{2} \left(|q|^2 + \frac{1}{|q|^2} \right) + \frac{1}{2} |\rho|^2 \right] = |h|^2 \left[\cosh(2\beta_{\mathbb{I}}) + \frac{1}{2} |\rho|^2 \right] = g_{YM}^2 \quad (20)$$

This implies that when $\rho \neq 0$ the coupling constant h at the conformal fixed point will be different from g_{YM} , in contrast to what happens for $\beta = \beta_{\mathbb{R}} \in \mathbb{R}$. In this sense, turning on ρ is similar to turning on the imaginary part of $\beta = \beta_{\mathbb{I}}$. Yet, there seems to exist a particular point in the deformation space for which $h = g_{YM}$ continues to hold in the large N limit. This occurs when:

$$\cosh(2\beta_{\mathbb{I}}) + \frac{1}{2} |\rho|^2 = 1 \Rightarrow \beta_{\mathbb{I}} = \frac{1}{2} \arg \cosh \left(1 - \frac{|\rho|^2}{2} \right) \quad (21)$$

Closer inspection however of (21) reveals that it has no possible solutions, assuming $\beta_{\mathbb{I}} \in \mathbb{R}$ and $|\rho| > 0$. This implies that despite appearances, there is no special point for which $h = g_{YM}$ at two loops in the planar limit. Naturally, one expects that an analogous equation relating the two couplings, for which $h = g_{YM}$ at large N , may arise at any order in perturbation theory. What is not clear of course, is whether it will generically have any solutions or not.

3 Marginal deformations and Noncommutativity

In [20] we showed that for the β -deformed gauge theory it is possible to construct a noncommutativity matrix Θ encoding in a precise manner information on the moduli space of the theory. This construction is very simple and is based on fundamental properties of the gauge theory and AdS/CFT. In what follows we will adopt the reasoning outlined in [20] in order to determine a noncommutativity matrix for the ρ -deformation. We set $\beta = 0$ for the time being and later on discuss how to incorporate $\beta \neq 0$.

Our starting point is the F-term constraints:

$$\begin{aligned} \Phi_1 \Phi_2 &= \Phi_2 \Phi_1 + \rho \Phi_3^2, & \Phi_2 \Phi_3 &= \Phi_3 \Phi_2 + \rho \Phi_1^2, & \Phi_3 \Phi_1 &= \Phi_1 \Phi_3 + \rho \Phi_2^2 \\ \bar{\Phi}_1 \bar{\Phi}_2 &= \bar{\Phi}_2 \bar{\Phi}_1 - \bar{\rho} \bar{\Phi}_3^2, & \bar{\Phi}_2 \bar{\Phi}_3 &= \bar{\Phi}_3 \bar{\Phi}_2 - \bar{\rho} \bar{\Phi}_1^2, & \bar{\Phi}_3 \bar{\Phi}_1 &= \bar{\Phi}_1 \bar{\Phi}_3 - \bar{\rho} \bar{\Phi}_2^2 \end{aligned} \quad (22)$$

from which we read the holomorphic and antiholomorphic parts of Θ interpreting the eigenvalues of these matrices in the large N limit, as noncommuting coordinates parametrizing the space transverse to the worldvolume of the D3-brane. More precisely we have:

$$\begin{aligned} [z_1, z_2] &= \rho z_3^2, & [z_2, z_3] &= \rho z_1^2, & [z_3, z_1] &= \rho z_2^2 \\ [\bar{z}_1, \bar{z}_2] &= -\bar{\rho} \bar{z}_3^2, & [\bar{z}_2, \bar{z}_3] &= -\bar{\rho} \bar{z}_1^2, & [\bar{z}_3, \bar{z}_1] &= -\bar{\rho} \bar{z}_2^2 \end{aligned} \quad (23)$$

Following [20] we would like to assume that there exists a star product between some *commuting* variables $z^I, \bar{z}^{\bar{I}}$ which leads to commutation relations analogous to (23), so that we can write for instance: $i\Theta^{12} = [z_1, z_2]_* = z_1 * z_2 - z_2 * z_1 = \rho z_3^2$. This enables us to define a noncommutativity matrix which although position dependent, its entries are ordinary commuting objects. Then, under a change of coordinates Θ will

transform as a contravariant antisymmetric tensor field. We therefore write Θ in matrix form as:

$$\Theta = \begin{pmatrix} 0 & i\rho z_3^2 & -i\rho z_2^2 & & & \\ -i\rho z_3^2 & 0 & i\rho z_1^2 & & ? & \\ i\rho z_2^2 & -i\rho z_1^2 & 0 & & & \\ & ? & & 0 & -i\bar{\rho}\bar{z}_3^2 & i\bar{\rho}\bar{z}_2^2 \\ & & & i\bar{\rho}\bar{z}_3^2 & 0 & -i\bar{\rho}\bar{z}_1^2 \\ & & & -i\bar{\rho}\bar{z}_2^2 & i\bar{\rho}\bar{z}_1^2 & 0 \end{pmatrix} \quad (24)$$

It is clear from (24) that the F-term constraints determine the (2,0) and (0,2) parts of Θ . D-terms will in principle specify the (1,1) pieces of the noncommutativity matrix. However, as demonstrated in [20], there is an alternative indirect way of acquiring the information pertaining to D-terms. Recall that for the β -deformed gauge theory it was possible to fully determine Θ by imposing certain simple conditions on its form — namely definite reality properties, symmetries and marginality. If and only if there exists a choice for the Θ^{IJ} components of the noncommutativity matrix and the parameter ρ which respects these requirements, can we hope to describe the deformation in noncommutative terms.⁴ We will see in the following that this is indeed the case here.

Let us first find out what are the possible (1,1) pieces of Θ which respect the symmetries of the theory. Consider for instance the commutator $[z^1, \bar{z}^2] = i\Theta^{1\bar{2}}(z, \bar{z})$. We easily see that: $[z^1, \bar{z}^2] \xrightarrow{\mathbb{Z}_{3(2)}} e^{-\frac{i2\pi}{3}} [z^1, \bar{z}^2]$. This constrains $\Theta^{1\bar{2}}$ to either vanish or be a combination of any of the following: $\bar{z}^1 z^3, \bar{z}^3 z^2, z^1 \bar{z}^2$. Note that all of the choices displayed are also invariant under the other discrete symmetry of the theory $\mathbb{Z}_{3(1)}$ as they should. In a similar fashion, one can determine all the other possible components of $\Theta^{I\bar{J}}$. Overall, this yields a plethora of potential noncommutativity parameters. Transforming however Θ to spherical coordinates⁵ and requiring that it be real, transverse to and independent of the radial direction r , uniquely determines Θ . To be more precise, there remain *two* different possibilities for Θ^{IJ} . One of them is valid for $\rho \equiv -q_1 \in \mathbb{R}$:

$$\Theta_1 = iq_1 \begin{pmatrix} 0 & z_3^2 & -z_2^2 & 0 & -z_3\bar{z}_1 + z_2\bar{z}_3 & z_2\bar{z}_1 - z_3\bar{z}_2 \\ -z_3^2 & 0 & z_1^2 & \bar{z}_2 z_3 - z_1\bar{z}_3 & 0 & -z_1\bar{z}_2 + z_3\bar{z}_1 \\ z_2^2 & -z_1^2 & 0 & -z_2\bar{z}_3 + z_1\bar{z}_2 & z_1\bar{z}_3 - z_2\bar{z}_1 & 0 \\ 0 & -z_3\bar{z}_2 + z_1\bar{z}_3 & z_2\bar{z}_3 - z_1\bar{z}_2 & 0 & -\bar{z}_3^2 & \bar{z}_2^2 \\ z_3\bar{z}_1 - z_2\bar{z}_3 & 0 & -z_1\bar{z}_3 + z_2\bar{z}_1 & \bar{z}_3^2 & 0 & -\bar{z}_1^2 \\ -z_2\bar{z}_1 + z_3\bar{z}_2 & z_1\bar{z}_2 - z_3\bar{z}_1 & 0 & -\bar{z}_2^2 & \bar{z}_1^2 & 0 \end{pmatrix} \quad (25)$$

and the other one, for $\rho \equiv iq_2$ with $q_2 \in \mathbb{R}$:

$$\Theta_2 = q_2 \begin{pmatrix} 0 & z_3^2 & -z_2^2 & 0 & z_3\bar{z}_1 + z_2\bar{z}_3 & -z_2\bar{z}_1 - z_3\bar{z}_2 \\ -z_3^2 & 0 & z_1^2 & -\bar{z}_2 z_3 - z_1\bar{z}_3 & 0 & z_1\bar{z}_2 + z_3\bar{z}_1 \\ z_2^2 & -z_1^2 & 0 & z_2\bar{z}_3 + z_1\bar{z}_2 & -z_1\bar{z}_3 - z_2\bar{z}_1 & 0 \\ 0 & z_3\bar{z}_2 + z_1\bar{z}_3 & -z_2\bar{z}_3 - z_1\bar{z}_2 & 0 & \bar{z}_3^2 & -\bar{z}_2^2 \\ -z_3\bar{z}_1 - z_2\bar{z}_3 & 0 & z_1\bar{z}_3 + z_2\bar{z}_1 & -\bar{z}_3^2 & 0 & \bar{z}_1^2 \\ z_2\bar{z}_1 + z_3\bar{z}_2 & -z_1\bar{z}_2 - z_3\bar{z}_1 & 0 & \bar{z}_2^2 & -\bar{z}_1^2 & 0 \end{pmatrix} \quad (26)$$

Combining the two into $\Theta_\rho = \Theta_1 + \Theta_2$ we define a unique noncommutativity matrix Θ describing the ρ -deformation for general complex $\rho = (-q_1 + iq_2) \in \mathbb{C}$. This presumably indicates that a noncommutative description of the transverse space is valid throughout the whole of the ρ parameter space, contrary to what happens for the β -deformed gauge theory.

Let us now examine Θ in order to determine its properties. Recall that the noncommutativity parameter for the β -deformed theory, turned out to be position independent along isometry directions of the metric. This was crucial for employing the Lunin-Maldacena generating technique. In this case, we obviously do not expect Θ to be constant along isometry directions since we know that the ρ -deformed theory does not respect

⁴Note for instance, that this description was not valid for the β -deformed theory when $\beta \in \mathbb{I}$.

⁵Refer to appendix A for the noncommutativity matrix in different coordinate systems.

any other global $U(1)$ symmetries except for the R -symmetry. Indeed, Θ is of a highly nontrivial form even when written in spherical coordinates (See appendix A). We may however hope to find a coordinate system for which Θ is position independent, even if not along isometry directions ⁶. In [20] we presented what we believe are the two necessary and sufficient conditions for this to occur:

$$\left. \begin{array}{l} \text{Divergence Free Condition} \\ \text{Associativity Condition} \end{array} \right\} \Rightarrow T^{[ijk]} = \partial_l(\Theta^{l[i}\Theta^{jk]}) = 0 \quad (27)$$

$$T^{[ijk]} \equiv \Theta^{il}\partial_l\Theta^{jk} + \Theta^{kl}\partial_l\Theta^{ij} + \Theta^{jl}\partial_l\Theta^{ki} = 0$$

It is actually easy to see that although the first condition is satisfied by the noncommutativity matrix thus determined, the second one is not. This implies that contrary to the β -deformation, there doesn't exist a coordinate system in which Θ can be put in constant form ⁷. More importantly however, nonassociativity makes the task of explicitly constructing an appropriate star product for the scalar fields a rather non-trivial one. In fact, we have not so far been able to find a star product so as to rewrite the Lagrangian of the ρ -deformed gauge theory as that of the $\mathcal{N} = 4$ Lagrangian with the usual product between the matter content of the theory replaced by the star product. We will further address this issue in relation to the applicability of the method used in this article in section 7.

Finally, note that failure of associativity stems particularly from the (1,1) parts of the noncommutativity matrix thus naturally challenging our method for determining them. There exists however what we believe to be a highly non-trivial check that we have constructed the correct Θ describing the deformation. We saw in the previous section, that for some special points in the space of couplings of the marginally deformed theory, one can move from a theory where either β or ρ (but not both) is turned on, to a theory where both couplings are nonvanishing. The whole analysis as well as the appropriate field redefinitions which took us from one point to the other in the deformation space, relied on the holomorphicity of the superpotential. It would thus appear quite improbable that we would be able to see it happening in this context. In principle however, one would expect that if the deformation is indeed described from an open string theory perspective as a noncommutative deformation of the transverse space, then at these special points Θ should transform under a change of coordinates from Θ_β or Θ_ρ to $\Theta = \Theta_{\tilde{\beta}} + \Theta_{\tilde{\rho}}$. Moreover, one might hope that the coordinate transformation which would make this possible would be the precise analog of the field redefinition applied to the gauge theory. Note however that in the case of the β -deformation, it is only for $\beta = \beta_{\mathbb{R}} \in \mathbb{R}$ that a noncommutative description — with parameter $a = 2\sin\beta_{\mathbb{R}}$ — is valid. This implies that we can apply the above consistency check if and only if both the original and final points in the coupling constant space involve a real parameter $\beta_{\mathbb{R}}$. A glance at the previous section will convince us that this indeed occurs: starting with $\rho = q_1 \in \mathbb{R}$ and $\beta = 0$ one can reach a point with $\tilde{\rho} = \frac{iq_1}{\sqrt{3}} \in \mathbb{I}$ and $\tilde{a} = 2\sin\tilde{\beta} = \frac{2q_1}{\sqrt{3}} \in \mathbb{R}$. In fact it is quite straightforward to check that a coordinate transformation according to (19) leads us from $\Theta_\rho = -\Theta_1$ to $\Theta = \Theta_{\tilde{a}=\frac{2q_1}{\sqrt{3}}} + \Theta_{\tilde{\rho}=\frac{iq_1}{\sqrt{3}}}$. Furthermore, it appears that this case exhausts all possible coordinate changes that relate noncommutativity matrices corresponding to different parameters of the Leigh–Strassler deformation. We take this result as evidence that both our prescription for determining the (1,1) parts of Θ as well as the very interpretation of the deformation in noncommutative terms are indeed justified.

4 The Seiberg–Witten equations and the deformed flat space solution.

In the previous section, we saw how the deformation of the superpotential affects the moduli space of the gauge theory at large N . In particular, the six dimensional flat space with metric $\mathcal{G}_{IJ}$ of the $\mathcal{N} = 4$ theory is promoted to a noncommutative space characterized now by the set $\mathcal{G}_{IJ}$ and Θ^{IJ} . Both metric and noncommutativity parameter are mainly determined from the Lagrangian of the theory; the former is read off from the kinetic term of the scalars while the latter from their potential.

⁶This is the case for the nongeometric Q -space [82, 83, 84], for instance.

⁷Note that this does not of course exclude the possibility of finding a reference frame for which this is true. Integrability however will be lost.

Since an $SU(N)$ gauge theory can be realized as the low energy limit of open strings attached on a stack of D3-branes, the set $(\mathcal{G}_{flat}, \Theta)$ describes the geometry of the transverse space as seen by the open strings in the limit of large N and $\alpha' \rightarrow 0$. We will thus refer to $(\mathcal{G}_{flat}, \Theta)$ as the *open* string parameters.

On the other hand, any theory of open strings necessarily contains closed strings. Closed strings however perceive the geometry quite differently from open strings. In fact, it was shown in [21, 22] that target space noncommutativity from the point of view of open strings corresponds to turning on a B-field from the viewpoint of closed strings. The set (g, B) , with g the closed string metric, are the *closed* string parameters that describe the same geometry. In this context, (g, B) represent the deformed flat space solution into which D3-branes are immersed⁸. Suppose now that we are given a set of equations relating the two groups of data. Then — provided that the open string parameters determined in the previous section exactly and fully describe the deformation — we could specify the closed string fields (g, B) of the deformed flat space geometry for free, i.e. without having to solve the type IIB differential equations of motion [85].

Equations relating open and closed string parameters indeed exist in the literature [86, 87, 22, 21]:

$$\begin{aligned} g + B &= \frac{1}{\mathcal{G}^{-1} + \Theta} \\ g_s &= G_s \sqrt{\frac{\det \mathcal{G}^{-1}}{\det (\mathcal{G}^{-1} + \Theta)}} = G_s \sqrt{\frac{1}{\det (1 + \Theta \mathcal{G})}} \end{aligned} \quad (28)$$

where G_s, g_s denote the corresponding open/closed string couplings⁹. They were however considered in a situation somewhat different from the one discussed in this article, namely for a *flat* D-brane embedded in *flat* background space with a *constant* B-field turned on *along* its worldvolume [22, 21]. It was under these circumstances that, the presence of the background B-field was shown to deform the algebra of functions on the worldvolume of the brane into that of a noncommutative Moyal type of algebra, where Θ is a c-number. While it is natural to ask what happens in situations where the B-field is not constant, technical difficulties have hindered progress in this direction. In the order of increasing complexity, two cases can be considered: the case of a closed $dB = 0$ though not necessarily constant two-form field B and the case of nonvanishing NS-NS three form flux $H = dB$ in a curved background. In [88] the former case was explored and the Moyal deformation of the algebra of functions on the brane worldvolume was shown to naturally extended to the Kontsevich star product deformation [89]. The authors of [90] — see also [91, 92, 93] — undertook the study of the most general case where $H = dB \neq 0$. They considered a special class of closed string backgrounds, called parallelizable, and expanded the background fields in Taylor series. It was then possible to perturbatively analyze n-point string amplitudes on the disk and obtain — in a first order expansion — the appropriate generalization of (28). In fact, it turned out that equation (28) is still valid for a weakly varying nonclosed B-field even though the corresponding algebra of functions is now both noncommutative and nonassociative.

In this letter, we want to apply the above formulas in a situation where the B-field lies in the *transverse* space to the D3-brane. Despite the fact that this case has not been explicitly studied in the literature¹⁰ one expects by T-duality that equations (28) should continue to hold. If our reasoning thus far is correct and (28) indeed provide the relation between open and closed string parameters in this setup, the resulting closed string fields (g_s, g, B) will constitute a new supergravity solution, i.e. the deformed flat space solution where D3-branes embedded.

There exists a natural place where we can test these ideas prior to checking whether the type IIB field equations for the set (g_s, g, B) are satisfied. Recall that both the gravity dual and the corresponding deformed flat space background are known for the β -deformed gauge theory [19]. The open string data $(\mathcal{G}_{flat}, \Theta_{\beta_R})$ describing the β -deformation were determined in [20] where the noncommutativity parameter was found to be constant despite the fact that the associated NS-NS three form flux was non zero. It is easy to show that applying (28) to the open string parameters $(\mathcal{G}_{flat}, \Theta_{\beta_R})$ one recovers the deformed flat space geometry found by Lunin and Maldacena in [19]. This may appear as a surprise unless one observes the extraordinary

⁸We are obviously interested here in the limit where open and closed strings are decoupled from each other.

⁹Note that $G_s = 1$ for the ρ -deformation.

¹⁰Mainly because a constant B-field in the transverse space can be gauged away leaving no trace on the geometry.

similarity between (28) and the T-duality transformation rules of (8). In fact, these equations are identical in this case although the interpretation of the variables involved is essentially different. We will return to this point again in the following section.

Having tested our ideas in the context of the β -deformation, we proceed to check whether one can specify the appropriate background for the ρ -deformation as well. It actually turns out that the closed string data (g_s, g, B) determined in the fashion described above, satisfy the supergravity equations of motion only up to *third* order in the deformation parameter ρ . The result is at least perplexing — whereas it doesn't completely invalidate our considerations it indicates the presence of some kind of flow in them. In fact, the discrepancy at higher orders can be traced in a variety of reasons. Note however that the breakdown of (28) in the case of a nonassociative deformation seems to be the most plausible one, since nonassociativity manifests itself at second order in the deformation parameter. We postpone further discussion on this issue until section 6 and close this section by presenting the detailed — though not particularly illuminating — form of the solution to this order ¹¹.

Metric g :

$$\begin{aligned}
g_{r_1 r_1} &= 1 - (r_2^2 [(q_1 x_3 - q_2 y_1)^2 + (q_1 y - q_2 x_2)^2] + r_3^2 [(q_1 y - q_2 x_3)^2 + (q_1 x_2 - q_2 y_1)^2]) \\
g_{r_2 r_2} &= 1 - (r_1^2 [(q_1 x_3 - q_2 y_2)^2 + (q_1 y - q_2 x_1)^2] + r_3^2 [(q_1 y - q_2 x_3)^2 + (q_1 x_1 - q_2 y_2)^2]) \\
g_{r_3 r_3} &= 1 - (r_1^2 [(q_1 x_2 - q_2 y_3)^2 + (q_1 y - q_2 x_1)^2] + r_2^2 [(q_1 y - q_2 x_2)^2 + (q_1 x_1 - q_2 y_3)^2]) \\
g_{\varphi_1 \varphi_1} &= r_1^2 [1 - (r_2^2 [(q_1 x_1 - q_2 y_3)^2 + (q_1 y_2 - q_2 x)^2] + r_3^2 [(q_1 y_3 - q_2 x)^2 + (q_1 x_1 - q_2 y_2)^2])] \\
g_{\varphi_2 \varphi_2} &= r_2^2 [1 - (r_1^2 [(q_1 x_2 - q_2 y_3)^2 + (q_1 y_1 - q_2 x)^2] + r_3^2 [(q_1 y_3 - q_2 x)^2 + (q_1 x_2 - q_2 y_1)^2])] \\
g_{\varphi_3 \varphi_3} &= r_3^2 [1 - (r_1^2 [(q_1 x_3 - q_2 y_2)^2 + (q_1 y_1 - q_2 x)^2] + r_2^2 [(q_1 y_2 - q_2 x)^2 + (q_1 x_3 - q_2 y_1)^2])] \\
g_{r_1 r_2} &= r_1 r_2 [(q_1 x_3 - q_2 y_1)(q_1 x_3 - q_2 y_2) + (q_1 y - q_2 x_1)(q_1 y - q_2 x_2)] \\
g_{r_1 r_3} &= r_1 r_3 [(q_1 x_2 - q_2 y_1)(q_1 x_2 - q_2 y_3) + (q_1 y - q_2 x_3)(q_1 y - q_2 x_1)] \\
g_{r_1 \varphi_1} &= r_1 (r_2^2 [(q_1 x_3 - q_2 y_1)(q_2 x - q_1 y_2) + (q_2 x_2 - q_1 y)(q_1 x_1 - q_2 y_3)] + \\
&\quad + r_3^2 [(q_2 x_3 - q_1 y)(q_1 x_1 - q_2 y_2) + (q_1 x_2 - q_2 y_1)(q_2 x - q_1 y_3)]) \\
g_{r_1 \varphi_2} &= r_1 r_2^2 [(q_2 x - q_1 y_1)(-q_1 x_3 + q_2 y_1) + (q_2 x_2 - q_1 y)(-q_1 x_2 + q_2 y_3)] \\
g_{r_1 \varphi_3} &= r_1 r_3^2 [(q_2 x - q_1 y_1)(-q_1 x_2 + q_2 y_1) + (q_2 x_3 - q_1 y)(-q_1 x_3 + q_2 y_2)] \\
g_{r_2 r_3} &= r_2 r_3 [(q_1 x_1 - q_2 y_3)(q_1 x_1 - q_2 y_2) + (q_1 y - q_2 x_3)(q_1 y - q_2 x_2)] \\
g_{r_2 \varphi_1} &= r_2 r_1^2 [(q_2 x - q_1 y_2)(-q_1 x_3 + q_2 y_2) + (q_2 x_1 - q_1 y)(-q_1 x_1 + q_2 y_3)] \\
g_{r_2 \varphi_2} &= r_2 (r_1^2 [(q_1 x_3 - q_2 y_2)(q_2 x - q_1 y_1) + (q_2 x_1 - q_1 y)(q_1 x_2 - q_2 y_3)] + \\
&\quad + r_3^2 [(q_2 x_3 - q_1 y)(q_1 x_2 - q_2 y_1) + (q_1 x_1 - q_2 y_2)(q_2 x - q_1 y_3)]) \\
g_{r_2 \varphi_3} &= r_2 r_3^2 [(q_2 x_3 - q_1 y)(-q_1 x_3 + q_2 y_1) + (q_2 x - q_1 y_2)(-q_1 x_1 + q_2 y_2)] \\
g_{r_3 \varphi_1} &= r_3 r_1^2 [(q_2 x_1 - q_1 y)(-q_1 x_1 + q_2 y_2) + (q_2 x - q_1 y_3)(-q_1 x_2 + q_2 y_3)] \\
g_{r_3 \varphi_2} &= r_3 r_2^2 [(q_2 x_2 - q_1 y)(-q_1 x_2 + q_2 y_1) + (q_2 x - q_1 y_3)(-q_1 x_1 + q_2 y_3)] \\
g_{r_3 \varphi_3} &= r_3 (r_2^2 [(q_1 x_3 - q_2 y_1)(q_2 x_2 - q_1 y) + (q_2 x - q_1 y_2)(q_1 x_1 - q_2 y_3)] + \\
&\quad + r_1^2 [(q_2 x_1 - q_1 y)(q_1 x_3 - q_2 y_2) + (q_1 x_2 - q_2 y_3)(q_2 x - q_1 y_1)]) \\
g_{\varphi_1 \varphi_2} &= r_1^2 r_2^2 [(q_2 x - q_1 y_1)(-q_1 x + q_2 x) + (q_2 y_3 - q_1 x_2)(-q_1 x_1 + q_2 y_3)] \\
g_{\varphi_1 \varphi_3} &= r_1^2 r_3^2 [(q_2 x - q_1 y_1)(-q_1 y_3 + q_2 x) + (q_2 y_2 - q_1 x_3)(-q_1 x_1 + q_2 y_2)] \\
g_{\varphi_2 \varphi_3} &= r_2^2 r_3^2 [(q_2 x - q_1 y_2)(-q_1 y_3 + q_2 x) + (q_2 y_1 - q_1 x_3)(-q_1 x_2 + q_2 y_1)]
\end{aligned} \tag{29}$$

¹¹ See appendix for the precise definition of the variables x, x_i, y, y_i which appear here.

Dilaton:

$$\begin{aligned}
e^{2\Phi} &= G \\
G &= 1 + r_1^2 [(q_1y - q_2x_1)^2 + (q_1y_1 - q_2x)^2 + (q_1x_3 - q_2y_2)^2 + (q_1x_2 - q_2y_3)^2] + \\
&\quad + r_2^2 [(q_1y - q_2x_2)^2 + (q_1y_2 - q_2x)^2 + (q_1x_1 - q_2y_3)^2 + (q_1x_3 - q_2y_1)^2] + \\
&\quad + r_3^2 [(q_1y - q_2x_3)^2 + (q_1x_2 - q_2y_1)^2 + (q_1x_1 - q_2y_2)^2 + (q_1y_3 - q_2x)^2]
\end{aligned} \tag{30}$$

B-field:

$$\begin{aligned}
B_{r_1r_2} &= r_3(q_2x_3 - q_1y) & B_{r_2r_3} &= r_1(q_2x_1 - q_1y) & B_{r_3r_1} &= r_2(q_2x_2 - q_1y) \\
B_{r_1\varphi_2} &= -r_2r_3(q_1x_2 - q_2y_1) & B_{r_1\varphi_3} &= r_2r_3(q_1x_3 - q_2y_1) & B_{r_2\varphi_1} &= r_1r_3(q_1x_1 - q_2y_2) \\
B_{r_2\varphi_3} &= -r_1r_3(q_1x_3 - q_2y_2) & B_{r_3\varphi_1} &= -r_1r_2(q_1x_1 - q_2y_3) & B_{r_3\varphi_2} &= r_1r_2(q_1x_2 - q_2y_3) \\
B_{\varphi_1\varphi_2} &= r_1r_2r_3(q_2x - q_1y_3) & B_{\varphi_2\varphi_3} &= r_1r_2r_3(q_2x - q_1y_1) & B_{\varphi_3\varphi_1} &= r_1r_2r_3(q_2x - q_1y_2)
\end{aligned} \tag{31}$$

5 D-branes in deformed $\text{AdS}_5 \times \text{S}^5$ and the near horizon geometry.

In this section we will address the issue of finding the gravity dual of the ρ -deformed gauge theory. To this end, it is helpful to first consider the β -deformation. As mentioned in the previous section, the T-duality transformation rules (8) with which the Lunin-Maldacena solution was constructed are identical in form to (28). Recall from section 2, that in order to obtain the dual background in this case one must use:

$$E_0 = g_{\text{AdS}_5 \times \text{S}^5} \quad \text{and} \quad \Gamma = \Theta_{\beta_{\mathbb{R}}} \tag{32}$$

Suppose now that we want to interpret these variables according to (28). We would obviously have to think of $g_{\text{AdS}_5 \times \text{S}^5}$ as the *open* string metric $\mathcal{G}_{\text{AdS}_5 \times \text{S}^5}$ whereas of Γ as $\Theta_{\beta_{\mathbb{R}}}$. In this sense, $(G_s = g_{YM}^2, \mathcal{G}_{\text{AdS}_5 \times \text{S}^5}, \Theta_{\beta_{\mathbb{R}}})$ would encode the geometry as seen at large N by the open strings attached on a D3-brane embedded in the Lunin-Maldacena (11) background. In the following we denote the NS-NS fields of the solution as $(\tilde{g}_s, \tilde{g}, \tilde{B})$.

In other words, consider a stack of N D3-branes in the deformed flat space geometry of (10). The near horizon limit of this configuration is the gravity dual of the Leigh-Strassler marginal deformation with $\beta = \beta_{\mathbb{R}} \in \mathbb{R}$ and $\rho = 0$. A probe D3-brane propagating near the stack will then be described by the DBI action written either in terms of the closed $(\tilde{g}_s, \tilde{g}, \tilde{B})$ or of the open $(G_s, \mathcal{G}_{\text{AdS}_5 \times \text{S}^5}, \Theta_{\beta})$ string fields. However, the action of a single D3-brane separated from a collection of (N-1) other branes can also be obtained by integrating out the massive open strings stretched between the probe and the source. Indeed, as expected according to [23, 24, 25, 26, 27, 28], the DBI action describing the motion of a D3-brane in this background should in the large N limit coincide with the leading IR part of the quantum effective action of the β -deformed theory obtained by keeping the U(1) external fields and integrating over the massive ones.

In this spirit, it does not seem surprising that the appropriate open string data in this case, are the metric of $\text{AdS}_5 \times \text{S}^5$ and the noncommutativity parameter $\Theta_{\beta_{\mathbb{R}}}$. In fact, the action of the β -deformed gauge theory can be written as that of the parent $\mathcal{N} = 4$ theory with the product of the matter fields replaced by a star product associated to $\Theta_{\beta_{\mathbb{R}}}$. Moreover, as conjectured in [19, 44, 45] and later proven in [29], all planar amplitudes are equal to their $\mathcal{N} = 4$ counterparts up to an overall phase factor. This suggests that the iterative structure of the large N β -deformed gauge theory amplitudes, when $\beta = \beta_{\mathbb{R}} \in \mathbb{R}$, is identical to that of the $\mathcal{N} = 4$ SYM theory. It is then not hard to imagine that the quantum effective action mentioned above will be analogous to that of the undeformed theory with the only difference being some phase factors coming from the noncommutative deformation of the product. Subsequently, the open string fields appearing in the DBI form of the effective action of the $\mathcal{N} = 4$ theory $(G_s, \mathcal{G}, \Theta = 0)$ will be promoted to $(G_s, \mathcal{G}, \Theta_{\beta_{\mathbb{R}}})$.

It is natural to wonder whether a similar situation could apply to the ρ -deformation as well. The analysis of section 3 may lead us to think that this is most likely *not* the case. Even if we succeeded in writing the action of the theory in question, as the $\mathcal{N} = 4$ action with a star product between the matter fields, it is difficult to understand how planar equivalence between the two would be achieved with the deformation being both noncommutative and nonassociative. In fact, the proof given in [29] specifically relied on the

associativity of the star product for the β -deformation. Nevertheless, nonassociativity is a second order effect in ρ and in view of the results of the previous section, one might hope that a solution to this order could be obtained in this case, too.

To explicitly check this we can directly use the second order expansion of (28):

$$\begin{aligned} g &= \mathcal{G} + \mathcal{G}\Theta\mathcal{G}\Theta\mathcal{G} + \mathcal{O}(\rho^4) \\ B &= -\mathcal{G}\Theta\mathcal{G} + \mathcal{O}(\rho^3) \\ G^{-1} &= 1 + \text{Tr} \left[\mathcal{G}\Theta - \frac{1}{2}\mathcal{G}\Theta\mathcal{G}\Theta \right] + \mathcal{O}(\rho^4) \end{aligned} \quad (33)$$

The equations above, provide a relation between the open string parameters of the deformed theory and the NS-NS string fields of the dual geometry. Since there is no obvious way to extract information on the associated RR-fluxes, we resort to the type IIB equations of motion in order to specify them. Fortunately, assuming no warp factor in front the metric and making the usual ansatz for the five form field strength ¹²:

$$\begin{aligned} ds_{10}^2 &= ds_{AdS_5}^2 + ds_{S^5}^2 \\ F_5 &= f(\omega_{AdS_5} + \omega_{S^5}) \end{aligned} \quad (34)$$

allows us to directly solve for the RR three form flux F_3 :

$$\begin{aligned} F_3 &= -f^{-1}d \star_5 e^{-2\Phi} H_3 \\ H_3 &= f^{-1}d \star_5 F_3 \Rightarrow d[B - f^{-1} \star_5 F_3] = 0 \end{aligned} \quad (35)$$

Note that to this order $F_3 = f \star_5 B$ which greatly simplifies calculations ¹³. One can then indeed show that the type IIB equations are simultaneously satisfied up to third order in the deformation parameter, for the following set of fields ¹⁴:

Dilaton:

$$\begin{aligned} e^{2\Phi} &= e^{2\Phi_0} G \\ G^{-1} &= 1 + q_1^2 (y^2 + s_\alpha^2 x_1^2 + (c_\alpha^2 + s_\alpha^2 c_\theta^2) x_2^2 + (c_\alpha^2 + s_\alpha^2 s_\theta^2) x_3^2 + c_\alpha^2 y_1^2 + s_\alpha^2 s_\theta^2 y_2^2 + s_\alpha^2 c_\theta^2 y_3^2) + \\ &\quad + 2q_1 q_2 (xy + (c_\alpha^2 - s_\alpha^2) x_1 y_1 - (c_\theta^2 + s_\theta^2 (c_\alpha^2 - s_\alpha^2)) x_2 y_2 - (s_\theta^2 + c_\theta^2 (c_\alpha^2 - s_\alpha^2))) \\ &\quad + q_1^2 (x^2 + c_\alpha^2 x_1^2 + s_\alpha^2 s_\theta^2 x_2^2 + s_\alpha^2 c_\theta^2 x_3^2 + s_\alpha^2 y_1^2 + (c_\alpha^2 + s_\alpha^2 c_\theta^2) y_2^2 + (c_\alpha^2 + s_\alpha^2 s_\theta^2) y_3^2) \end{aligned} \quad (36)$$

B-field:

$$\begin{aligned} B_{\alpha\theta} &= s_\alpha (q_1 y + q_2 x) \quad B_{\alpha\varphi_1} = 0 \quad B_{\alpha\varphi_2} = s_\alpha s_\theta c_\theta (q_1 x_2 - q_2 y_2) \quad B_{\alpha\varphi_3} = s_\alpha s_\theta c_\theta (-q_1 x_3 + q_2 y_3) \\ B_{\theta\varphi_1} &= c_\alpha s_\alpha^2 (q_1 x_1 - q_2 y_1) \quad B_{\theta\varphi_2} = -c_\alpha s_\alpha^2 s_\theta^2 (q_1 x_2 - q_2 y_2) \quad B_{\theta\varphi_3} = -c_\alpha s_\alpha^2 c_\theta^2 (q_1 x_3 - q_2 y_3) \\ B_{\varphi_1\varphi_2} &= -c_\alpha s_\alpha^2 s_\theta c_\theta (q_1 y_3 + q_2 x_3) \quad B_{\varphi_2\varphi_3} = -c_\alpha s_\alpha^2 s_\theta c_\theta (q_1 y_1 + q_2 x_1) \quad B_{\varphi_3\varphi_1} = -c_\alpha s_\alpha^2 s_\theta c_\theta (q_1 y_2 + q_2 x_2) \end{aligned} \quad (37)$$

F_3 and F_5 -form flux:

$$\begin{aligned} F_3 &= \star_{S^5} \Omega \quad \text{with} \quad \Omega \equiv \mathcal{G}_{ik} \mathcal{G}_{jl} \Theta^{kl} dx^i \wedge dx^j \\ F_5 &= f(\omega_{AdS_5} + G\omega_{S^5}) \end{aligned} \quad (38)$$

¹²We refer the reader to the appendix for the necessary definitions of the parameters involved as well as the type IIB field equations [85] in five dimensions.

¹³This is intimately connected to the fact that $\delta_{S^5} \Omega_\rho = 0$ where $\Omega_\rho = \mathcal{G}\Theta\mathcal{G}$ denotes the form on S^5 associated to the bivector Θ_ρ . It is clear from (33) that $B = -\Omega$ to this order. It is worth remarking that $F_3 = f \star B$ is exact for the β -deformed theory where both $d\Omega_\beta = 0$ and $\delta_{S^5} \Omega_\beta = 0$ hold.

¹⁴We set $R = 1$ where R the radius of $\text{Admathrm}S_5$.

Metric g :

$$\begin{aligned}
g_{\alpha\alpha} &= 1 - q_1^2 (c_\theta^2 x_2^2 + s_\theta^2 x_3^2 + y^2) + 2q_1 q_2 (-xy + c_\theta^2 x_2 y_2 + s_\theta^2 x_3 y_3) - q_2^2 (x^2 + c_\theta^2 y_2^2 + s_\theta^2 y_3^2) \\
g_{\theta\theta} &= s_\alpha^2 [1 - q_1^2 (y^2 + s_\alpha^2 x_1^2 + c_\alpha^2 (s_\theta^2 x_2^2 + c_\theta^2 x_3^2)) - 2q_1 q_2 (xy - s_\alpha^2 x_1 y_1 - c_\alpha^2 (s_\theta^2 x_2 y_2 + c_\theta^2 x_3 y_3)) - \\
&\quad - q_2^2 (x^2 + s_\alpha^2 y_1^2 + c_\alpha^2 (s_\theta^2 y_2^2 + c_\theta^2 y_3^2))] \\
g_{\varphi_1 \varphi_1} &= c_\alpha^2 [1 - s_\alpha^2 (q_1^2 (x_1^2 + s_\theta^2 y_2^2 + c_\theta^2 y_3^2) + 2q_1 q_2 (-x_1 y_1 + s_\theta^2 x_2 y_2 + c_\theta^2 x_3 y_3) + q_2^2 (y_1^2 + s_\theta^2 x_2^2 + c_\theta^2 x_3^2))] \\
g_{\varphi_2 \varphi_2} &= s_\alpha^2 s_\theta^2 [1 - q_1^2 (x_2^2 (c_\alpha^2 + s_\alpha^2 c_\theta^2) + c_\alpha^2 y_1^2 + s_\alpha^2 c_\theta^2 y_3^2) + 2q_1 q_2 (-c_\alpha^2 x_1 y_1 + (c_\alpha^2 + s_\alpha^2 c_\theta^2) x_2 y_2 - s_\alpha^2 c_\theta^2 x_3 y_3) - \\
&\quad - q_2^2 (c_\alpha^2 x_1^2 + s_\alpha^2 c_\theta^2 x_3^2 + (c_\alpha^2 + s_\alpha^2 c_\theta^2) y_2^2)] \\
g_{\varphi_3 \varphi_3} &= s_\alpha^2 c_\theta^2 [1 - q_1^2 (c_\alpha^2 y_1^2 + s_\alpha^2 s_\theta^2 y_2^2 + (c_\alpha^2 + s_\alpha^2 s_\theta^2) x_2^2) + 2q_1 q_2 (-c_\alpha^2 x_1 y_1 - s_\alpha^2 s_\theta^2 x_2 y_2 + (c_\alpha^2 + s_\alpha^2 s_\theta^2) x_3 y_3) - \\
&\quad - q_2^2 (c_\alpha^2 x_1^2 + s_\alpha^2 s_\theta^2 x_2^2 + (c_\alpha^2 + s_\alpha^2 s_\theta^2) y_3^2)] \\
g_{\alpha\theta} &= c_\alpha s_\alpha c_\theta s_\theta [q_1^2 (x_2^2 - x_3^2) + 2q_1 q_2 (-x_2 y_2 + x_3 y_3) + q_2^2 (y_2^2 - y_3^2)] \\
g_{\alpha\varphi_1} &= c_\alpha s_\alpha [q_1^2 (x_1 y + s_\theta^2 x_3 y_2 c_\theta^2 x_2 y_3) + q_1 q_2 (x x_1 - y y_1 + x_2 x_3 - y_2 y_3) - q_2^2 (x y_1 + c_\theta^2 x_3 y_2 + s_\theta^2 x_2 y_3)] \\
g_{\alpha\varphi_2} &= c_\alpha s_\alpha s_\theta^2 [-q_1^2 (x_2 y + x_3 y_1) + q_1 q_2 (-x x_2 - x_1 x_3 + y y_2 + y_1 y_3) + q_2^2 (x y_2 + x_1 y_3)] \\
g_{\alpha\varphi_3} &= c_\alpha s_\alpha s_\theta^2 [-q_1^2 (x_2 y + x_3 y_1) + q_1 q_2 (-x x_2 - x_1 x_3 + y y_2 + y_1 y_3) + q_2^2 (x y_2 + x_1 y_3)] \\
g_{\theta\varphi_1} &= c_\alpha^2 s_\alpha^2 c_\theta s_\theta (q_1^2 + q_2^2) (x_3 y_2 - x_2 y_3) \\
g_{\theta\varphi_2} &= s_\alpha^2 s_\theta c_\theta [q_2^2 (x y_2 + s_\alpha^2 x_3 y_1 + c_\alpha^2 x_1 y_3) - q_1 q_2 (x x_2 - y y_2 + x_1 x_3 - y_1 y_3) - q_1^2 (x_1 y + c_\alpha^2 x_3 y_1 + s_\alpha^2 x_1 y_3)] \\
g_{\theta\varphi_3} &= s_\alpha^2 s_\theta c_\theta [q_1^2 (x_3 y + c_\alpha^2 x_2 y_1 + s_\alpha^2 x_1 y_2) + q_1 q_2 (x_1 x_2 - y_1 y_2 + x x_3 - y y_3) - q_2^2 (x y_3 + s_\alpha^2 x_2 y_1 + c_\alpha^2 x_1 y_2)] \\
g_{\varphi_1 \varphi_2} &= c_\alpha^2 s_\alpha^2 s_\theta^2 (q_1^2 + q_2^2) (x_1 x_2 + y_1 y_2) \\
g_{\varphi_2 \varphi_3} &= c_\alpha^2 s_\alpha^2 c_\theta^2 (q_1^2 + q_2^2) (x_1 x_3 + y_1 y_3) \\
g_{\varphi_3 \varphi_1} &= s_\alpha^4 c_\theta^2 s_\theta^2 (q_1^2 + q_2^2) (x_2 x_3 + y_2 y_3)
\end{aligned} \tag{39}$$

6 Discussion

In the previous sections we set forth some new ideas that helped us obtain new gravity solutions up to third order in the deformation parameter ρ . Moreover, we observed that the same method provided the exact supergravity backgrounds related to the β -deformed gauge theory. We view these results as evidence that supports the basic ideas of our proposal which can be nicely summarized in figure . In this section we would like to discuss its obscure points; the ones that possibly underlie its failure to provide the solution to all orders in the deformation parameter.

The method proposed in this note consists of three steps. Let us separately consider the issues that arise in each one. The starting point consists of determining the appropriate open string data. In this, we mainly use information from the gauge theory Lagrangian. Obviously, the fact that the noncommutativity matrix specified in this case, fails to preserve the property of associativity is quite displeasing. In particular, it is not at all clear how to define an appropriate star product. Subsequently, there is no obvious way in which one can rewrite the Lagrangian of the ρ -deformed theory in terms of the $\mathcal{N} = 4$ SYM Lagrangian with a modified product between the matter fields. In fact, most of the star products that we attempted to define produced extra terms in the action, to second and third order in the deformation parameter ρ . Interestingly enough though, all the additional terms were essentially of the same form as the ones coming from the β and ρ deformation themselves. We hope to explore this point further in the future.

A related issue is that of the D-terms. In [20] we showed that we can rewrite these terms in the $\mathcal{N} = 4$ theory as a sum of the F-terms with a potential term involving the commutator between holomorphic and antiholomorphic matter fields:

$$\text{Tr}[\Phi_I, \tilde{\Phi}^J][\Phi_J, \tilde{\Phi}^J] = \text{Tr}[\Phi_I, \Phi_J][\tilde{\Phi}^I, \tilde{\Phi}^J] + \text{Tr}[\Phi_I, \tilde{\Phi}^J][\Phi_J, \tilde{\Phi}^I] \tag{40}$$

It was then clear that should we wish to only deform the F-terms of the potential, we must appropriately




Figure 1:

alter the commutator: $[\Phi_I, \tilde{\Phi}^J]$. For the β -deformed gauge theory, the $(1,1)$ pieces of the noncommutativity matrix precisely ensured that the D-terms remained unaffected by the deformation according to (40). The lack of a star product in the case of the ρ -deformation however, makes it impossible to perform this consistency check.

The next step of the method proposed herein, consisted in mapping the open string parameters to the closed string ones. The precise mapping was formulated through the equations of (28) which as discussed in section 5, were derived under particularly different conditions than the ones considered in this paper. Their validity in this case is therefore naturally disputable, even more so, in view of the nonassociativity of Θ . We believe that this actually constitutes the most plausible reason for the failure of our proposal to produce the exact supergravity background whereas at the same time explains why the method works to this order where nonassociativity precisely comes into play. In fact, it seems that when T^{ijk} of (27) is nonvanishing, both Θ and $T = \Theta\partial\Theta$ are necessary for defining the deformation. A natural generalization of (28) would then relate $(\mathcal{G}, \Theta, T = \Theta\partial\Theta)$ to $(g, B, H = dB)$ and presumably provide the deformed flat space solution to all orders in the deformation.

Note that we do not necessarily maintain that it would directly solve the problem of finding the dual gravity background as well. It may very well be that nonassociativity spoils the planar equivalence between the $\mathcal{N} = 4$ theory and its deformation. This would obviously be reflected on the form of the quantum effective action and therefore of the DBI, making it difficult to determine the relevant open string data ¹⁵.

It is also worth remarking that (28) provides a relation only between the NS-NS fields of the open and closed string backgrounds. Information pertaining to the RR-fields is however essential, especially for determining the dual gravitational solution. In fact, in section 6 we had to rely on a particular ansatz for the metric and the five form flux in order to fully specify the background. In this light, it may seem plausible that a different ansatz — a warp factor in front of the AdS part of the metric, in particular — could grant us the solution to all orders in the deformation parameter. The presence of a non-trivial warp factor may actually be related to the deviation of the coupling constant h in (1) from its original g_{YM} value. Indeed, in the case of the β -deformation, a warp factor is absent from the solution when $\beta \in \mathbb{R}$ and $h = g_{YM}$, while it is not when $\beta_{\mathbb{I}} \neq 0$ and the Leigh-Strassler constraint indicates that $h \neq g_{YM}$. This fact therefore represents another possible explanation as to why our method fails to give an exact solution ¹⁶.

¹⁵Actually it is possible to imagine that this can occur independently from the nonassociativity of the product. Yet it would make it hard to understand why the procedure employed in this article would give a true solution to any order in the deformation.

¹⁶Note that we examined the case of a warp factor of the form G^n with $n \in \mathbb{R}$ and G of (36) but found no value of n for

Finally, we would like to note that throughout this article we considered the Leigh–Strassler marginal deformation at the point $\beta = 0$. It is however natural to think that quantum corrections will generically generate a β -like term since no obvious symmetry argument could prohibit it. In this sense it may seem rather significant to incorporate a nonvanishing β in our discussion. This is actually not difficult to do, provided that $\beta = \beta_{\mathbb{R}} \in \mathbb{R}$. In this case, we can define $\Theta = \Theta_{\beta_{\mathbb{R}}} + \Theta_{\rho}$ and follow the method outlined in this note. The result is straightforward but unfortunately does not give any further insight into the higher order corrections of the background. The case of generic $\beta \in \mathbb{C}$ is more interesting but also more difficult to study. A noncommutative description of the deformation is not valid in this case and one relies on the $SL(2, \mathbb{R})_s$ symmetry of the supergravity equations of motion in order to construct the dual solution [19]. Consequently, there is no obvious way to incorporate a complex β in our method.

The reason that makes the case of complex β worthwhile to explore further, is that according to the analysis of section 2, there exist some special points in the deformation space which can take us from a theory of generic β and $\rho = 0$, to a marginal deformation where both $\tilde{\rho}$ and $\tilde{\beta}$ are non vanishing. Since the gravity dual in the former case is known, investigating the solution at these points may provide useful information on how to extend our results to all orders in the deformation parameters.

7 Conclusion

In this article we studied the Leigh–Strassler marginal deformation of $\mathcal{N} = 4$ SYM for $\rho \neq 0$ and found gravity solutions corresponding to the associated flat space deformation and the AdS/CFT dual of the gauge theory, up to third order in the deformation parameter. We achieved this by first employing the prescription of [20] to obtain the open string parameters encoding the geometry of the moduli space and then determining the corresponding closed string fields.

There are various possibilities for future work that range from addressing the questions raised in the previous section, to establishing a precise connection with generalized complex geometry [94, 95], extending the relations between open and closed string parameters to include RR-fields — or rather generalizing the results of [96, 97, 98] to incorporate supersymmetry —, and finally apply similar considerations in the context of orbifold as well as relevant deformations of $\mathcal{N} = 4$ SYM and noncommutative gauge theories.

In summary, the ideas set forth in this note represent alternative means into investigating deformations of the AdS/CFT correspondence. Obviously, a number of issues should be resolved before they can provide a concrete proposal for constructing new supergravity backgrounds. We do however believe that they open up a path that leads to a better understanding of gauge/gravity duality which we therefore hope to further explore in the future.

Acknowledgments

I am indebted to A. Murugan and K. Zoubos for numerous discussions that greatly contributed to my understanding of several aspects of this work. I am also grateful to J. Maldacena and L. Rastelli for helpful conversations and encouragement as well as to J.P. Gauntlett, A. Naqvi, A. Parnachev, R. Ricci and M. Roček.

A The noncommutativity matrix

Here we present the noncommutativity matrix in polar coordinates (r_i, φ_i) with $i = 1, 2, 3$ on $\mathbb{R}^6$. We assume that Θ_{ρ} is given in terms of commuting variables $(z, \bar{z})$ and that we can follow the transformation rules of

which the supergravity equations were satisfied to third order in the deformation parameter.

contravariant tensors when changing coordinate systems, namely:

$$\Theta^{i'j'} = \frac{\partial x^{i'}}{\partial x^i} \frac{\partial x^{j'}}{\partial x^j} \Theta^{ij} \quad (41)$$

Rescaling q_i of (25) and (26) as $q_i \rightarrow 2q_i$ then yields:

$$\Theta_\rho = \begin{pmatrix} 0 & -(q_2 x_3 - q_1 y) r_3 & (q_2 x_2 - q_1 y) r_2 & 0 & \frac{(q_1 x_2 - q_2 y_1) r_3}{r_2} & \frac{(q_1 x_3 - q_2 y_1) r_2}{r_3} \\ (q_2 x_3 - q_1 y) r_3 & 0 & -(q_2 x_1 - q_1 y) r_1 & -\frac{(q_1 x_1 - q_2 y_2) r_3}{r_1} & 0 & \frac{(q_1 x_3 - q_2 y_2) r_1}{r_3} \\ -(q_2 x_2 - q_1 y) r_2 & (q_2 x_1 - q_1 y) r_2 & 0 & \frac{(q_1 x_1 - q_2 y_3) r_2}{r_1} & -\frac{(q_1 x_2 - q_2 y_3) r_1}{r_2} & 0 \\ 0 & \frac{(q_1 x_1 - q_2 y_2) r_3}{r_1} & -\frac{(q_1 x_1 - q_2 y_3) r_2}{r_1} & 0 & -\frac{(q_2 x - q_1 y_3) r_3}{r_1 r_2} & \frac{(q_2 x - q_1 y_2) r_2}{r_1 r_3} \\ -\frac{(q_1 x_2 - q_2 y_1) r_3}{r_2} & 0 & \frac{(q_1 x_2 - q_2 y_3) r_1}{r_2} & \frac{(q_2 x - q_1 y_3) r_3}{r_1 r_2} & 0 & -\frac{(q_2 x - q_1 y_1) r_1}{r_2 r_3} \\ \frac{(q_1 x_3 - q_2 y_1) r_2}{r_3} & -\frac{(q_1 x_3 - q_2 y_2) r_1}{r_3} & 0 & -\frac{(q_2 x - q_1 y_2) r_2}{r_1 r_3} & \frac{(q_2 x - q_1 y_1) r_1}{r_2 r_3} & 0 \end{pmatrix} \quad (42)$$

where to keep the expressions compact, we defined variables x, x_i and y, y_i according to:

$$\begin{aligned} x_1 &= -C_1 r_1 + C_2 r_2 + C_3 r_3 & x_2 &= C_1 r_1 - C_2 r_2 + C_3 r_3 & x_3 &= C_1 r_1 + C_2 r_2 - C_3 r_3 \\ y_1 &= -S_1 r_1 + S_2 r_2 + S_3 r_3 & y_2 &= S_1 r_1 - S_2 r_2 + S_3 r_3 & y_3 &= S_1 r_1 + S_2 r_2 - S_3 r_3 \\ x &= C_1 r_1 + C_2 r_2 + C_3 r_3 & y &= S_1 r_1 + S_2 r_2 + S_3 r_3 \end{aligned} \quad (43)$$

whereas S_i, C_i represent the following trigonometric functions:

$$\begin{aligned} S_1 &= \sin(\varphi_2 + \varphi_3 - 2\varphi_1), & S_2 &= \sin(\varphi_3 + \varphi_1 - 2\varphi_2), & S_3 &= \sin(\varphi_1 + \varphi_2 - 2\varphi_3) \\ C_1 &= \cos(\varphi_2 + \varphi_3 - 2\varphi_1), & C_2 &= \cos(\varphi_3 + \varphi_1 - 2\varphi_2), & C_3 &= \cos(\varphi_1 + \varphi_2 - 2\varphi_3) \end{aligned} \quad (44)$$

The discrete symmetry $\mathbb{Z}_{3(1)} \times \mathbb{Z}_{3(2)}$ along with the $U(1)_R$ are particularly transparent in this form. Observe first that under $\mathbb{Z}_{3(1)}$:

$$\begin{aligned} \mathbb{Z}_{3(1)} : \quad (x_1, x_2, x_3, y_1, y_2, y_3) &\rightarrow (x_3, x_1, x_2, y_3, y_1, y_2) \\ \text{while } (x, y) &\rightarrow (x, y) \end{aligned} \quad (45)$$

Then it is easy to see for example, that $\Theta_\rho^{r_1 \varphi_2} = \frac{(q_1 x_2 - q_2 y_1) r_3}{r_2} \rightarrow \Theta_\rho^{r_3 \varphi_1} = \frac{(q_1 x_1 - q_2 y_3) r_2}{r_1}$. The action of $\mathbb{Z}_{3(2)}$ is equally simple transforming the polar angles φ_i as:

$$\mathbb{Z}_{3(2)} : \quad (\varphi_1, \varphi_2, \varphi_3) \rightarrow (\varphi_1, \varphi_2 + \frac{2\pi}{3}, \varphi_3 - \frac{2\pi}{3}) \quad (46)$$

thus leaving invariant the trigonometric functions S_i, C_i which depend on the following combinations: $\sigma_i \equiv \frac{1}{3}(\varphi_{i+1} + \varphi_{i+2} - 2\varphi_1)$. Moreover, note that Θ_ρ is independent of $\psi = \frac{1}{3}(\varphi_1 + \varphi_2 + \varphi_3)$ therefore respects the $U(1)_R$ R-symmetry of the theory.

In a similar manner, one obtains the noncommutativity matrix Θ_ρ in spherical coordinates denoted as $(r, \alpha, \theta, \varphi_1, \varphi_2, \varphi_3)$.

$$\begin{aligned} z_1 &= r \cos \alpha e^{i\phi_1}, & z_2 &= r \sin \alpha \sin \theta e^{i\phi_2}, & z_3 &= r \sin \alpha \cos \theta e^{i\phi_3} \\ \bar{z}_1 &= r \cos \alpha e^{-i\phi_1}, & \bar{z}_2 &= r \sin \alpha \sin \theta e^{-i\phi_2}, & \bar{z}_3 &= r \sin \alpha \cos \theta e^{-i\phi_3} \end{aligned} \quad (47)$$

where it reads¹⁷:

$$\Theta_\rho = \begin{pmatrix} 0 & -\frac{q_2 x + q_1 y}{s_\alpha} & 0 & \frac{c_\theta(-q_1 x_2 + q_2 y_2)}{s_\alpha s_\theta} & \frac{s_\theta(q_1 x_3 - q_2 y_3)}{s_\alpha c_\theta} \\ \frac{q_2 x + q_1 y}{s_\alpha} & 0 & -\frac{q_1 x_1 + q_2 y_1}{c_\alpha} & \frac{c_\alpha(q_1 x_2 - q_2 y_2)}{s_\alpha^2} & \frac{c_\alpha(q_1 x_3 - q_2 y_3)}{s_\alpha^2} \\ 0 & \frac{q_1 x_1 - q_2 y_1}{c_\alpha} & 0 & \frac{c_\theta(q_2 x_3 + q_1 y_3)}{c_\alpha c_\theta} & -\frac{s_\theta(q_2 x_2 + q_1 y_2)}{c_\alpha c_\theta} \\ \frac{c_\theta(q_1 x_2 - q_2 y_2)}{s_\alpha s_\theta} & \frac{c_\alpha(-q_1 x_2 + q_2 y_2)}{s_\alpha^2} & -\frac{c_\theta(q_2 x_3 + q_1 y_3)}{c_\alpha c_\theta} & 0 & \frac{c_\alpha(q_2 x_1 + q_1 y_1)}{s_\alpha^2 s_\theta c_\theta} \\ -\frac{s_\theta(q_1 x_3 - q_2 y_3)}{s_\alpha c_\theta} & \frac{c_\alpha(-q_1 x_3 + q_2 y_3)}{s_\alpha^2} & \frac{s_\theta(q_2 x_2 + q_1 y_2)}{c_\alpha c_\theta} & -\frac{c_\alpha(q_2 x_1 + q_1 y_1)}{s_\alpha^2 s_\theta c_\theta} & 0 \end{pmatrix} \quad (48)$$

¹⁷We use here the following abbreviations: $s_\alpha = \sin \alpha, c_\alpha = \cos \alpha, s_\theta = \sin \theta, c_\theta = \cos \theta$.

Note that Θ_ρ is now a five-dimensional matrix along the S^5 and that variables x, x_i, y, y_i appearing in (48) are rescaled by $1/r$. In other words we have here defined:

$$\begin{aligned} x_1 &= (-c_\alpha C_1 + s_\alpha s_\theta C_2 + s_\alpha c_\theta C_3), & x_2 &= (c_\alpha C_1 - s_\alpha s_\theta C_2 + s_\alpha c_\theta C_3), & x_3 &= (c_\alpha C_1 + s_\alpha s_\theta C_2 - s_\alpha c_\theta C_3) \\ y_1 &= (-c_\alpha S_1 + s_\alpha s_\theta S_2 + s_\alpha c_\theta S_3), & y_2 &= (c_\alpha S_1 - s_\alpha s_\theta S_2 + s_\alpha c_\theta S_3), & y_3 &= (c_\alpha S_1 + s_\alpha s_\theta S_2 - s_\alpha c_\theta S_3) \\ y &= (c_\alpha S_1 + s_\alpha s_\theta S_2 + s_\alpha c_\theta S_3) & x &= (c_\alpha C_1 + s_\alpha s_\theta C_2 + s_\alpha c_\theta C_3) \end{aligned} \quad (49)$$

It is then clear that Θ_ρ is independent of the radial direction r .

B RR-fields and supergravity equations of motion

As mentioned previously, although the procedure proposed in this article gives us the solution for the NS-NS fields of the geometry for free, it does not produce any information on the RR-ones. We thus have to compute them using the supergravity equations of motions [85]. We employ the following ansatz¹⁸:

$$\begin{aligned} ds_{10}^2 &= ds_{AdS_5}^2 + ds_5^2 \\ C &= 0 \quad F_5 = f(\omega_{AdS_5} + \omega_{\tilde{S}^5}) \end{aligned} \quad (50)$$

where f is the appropriate normalization coefficient for the flux which in this case reduces to $f = 16\pi N$ and $\omega_{AdS_5}, \omega_{\tilde{S}^5}$ are the volume elements of the corresponding parts of the $AdS_5 \times \tilde{S}^5$ geometry. Then the supergravity field equations reduce to:

$$\begin{aligned} D^2 e^{-2\Phi} &= -\frac{1}{6} (F_3^2 - e^{-2\Phi} H_3^2) \\ F_3 &= -f^{-1} d \star_5 e^{-2\Phi} H_3 \\ H_3 &= f^{-1} d \star_5 F_3 \\ R_{MN} &= -2D_M D_N \Phi - \frac{1}{4} g_{MN} D^2 \Phi + \frac{1}{2} g_{MN} \partial_R \Phi \partial^R \Phi + \\ &\quad + \frac{1}{96} e^{2\Phi} F_{MPQR} F_N^{PQR} + \frac{1}{4} (H_{MPQ} H_N^{PQ} + e^{2\Phi} F_{MPQ} F_N^{PQ}) - \frac{1}{48} g_{MN} (H_3^2 + e^{2\Phi} F_3^2) \end{aligned} \quad (51)$$

where M, N represent five dimensional indices on the compact piece of the geometry whereas $\star_5$ denotes the Hodge star on the same manifold.

References

- [1] J. Maldacena, *The large N limit of Superconformal theories and supergravity*, *Adv. Theor. Math. Phys.* **2** (1998) 231, [[hep-th/9711200](#)].
- [2] S. S. Gubser, I. R. Klebanov, and A. M. Polyakov, *Gauge Theory Correlators from Non-Critical String Theory*, *Phys. Lett. B* **428** (1998) 105–114, [[hep-th/9802109](#)].
- [3] E. Witten, *Anti De Sitter Space And Holography*, *Adv. Theor. Math. Phys.* **2** (1998) 253–291, [[hep-th/9802150](#)].
- [4] I. Klebanov and E. Witten, *Superconformal Field Theory on Threebranes at a Calabi-Yau Singularity*, *Nucl. Phys. B* **536** (1998) 199–218, [[hep-th/9807080](#)].
- [5] I. R. Klebanov and M. Strassler, *Supergravity and a Confining Gauge Theory: Duality Cascades and χ SB-Resolution of Naked Singularities*, *JHEP* **0008** (2000) 052, [[hep-th/0007191](#)].

¹⁸Note that the vanishing axion condition can be deduced from the other two in equation (50).

- [6] J. Maldacena and C. Nunez, *Towards the large N limit of pure $\mathcal{N} = 1$ Super Yang Mills*, *Phys. Rev. Lett.* **86** (2001) 588–591, [[hep-th/0008001](#)].
- [7] S. Kachru and E. Silverstein, *4d Conformal Field Theories and Strings on Orbifolds*, *Phys. Rev. Lett.* **80** (1998) 4855–4858, [[hep-th/9802183](#)].
- [8] A. Lawrence, N. Nekrasov, and C. Vafa, *On Conformal Theories in Four Dimensions*, *Nucl. Phys. B* **533** (1998) 199–209, [[hep-th/9803015](#)].
- [9] J. P. Gauntlett, D. Martelli, and J. Sparks, *Supersymmetric AdS_5 solutions of M -theory*, *Class. Quant. Grav.* **21** (2004) 4335–4366, [[hep-th/0402153](#)].
- [10] K. Skenderis and M. Taylor, *Kaluza–Klein Holography*, *JHEP* **0605** (2006) 057, [[hep-th/0603016](#)].
- [11] K. Skenderis and M. Taylor, *Holographic Coulomb branch vevs*, *JHEP* **0608** (2006) 001, [[hep-th/0604169](#)].
- [12] J. P. Gauntlett, D. Martelli, J. Sparks, and D. Waldram, *Supersymmetric AdS Backgrounds in String and M -theory*, [hep-th/0411194](#).
- [13] J. P. Gauntlett, D. Martelli, J. Sparks, and D. Waldram, *Supersymmetric AdS_5 Solutions of Type IIB Supergravity*, *Class. Quant. Grav.* **23** (2006) 4693–4718, [[hep-th/0510125](#)].
- [14] J. Gauntlett, D. Martelli, J. Sparks, and S.-T. Yau, *Supersymmetric AdS_3 Solutions of Type IIB Supergravity*, [hep-th/0607080](#).
- [15] M. Wijnholt, *Parameter Space of Quiver Gauge Theories*, [hep-th/0512122](#).
- [16] K. Pilch and N. P. Warner, *A New Supersymmetric Compactification of Chiral IIB Supergravity*, *Phys. Lett. B* **487** (2000) 22–29, [[hep-th/0002192](#)].
- [17] K. Pilch and N. P. Warner, *$N=2$ Supersymmetric RG Flows and the IIB Dilaton*, *Nucl. Phys. B* **594** (2001) 209–228, [[hep-th/0004063](#)].
- [18] N. Halmagyi, K. Pilch, C. Romelsberger, and N. P. Warner, *Holographic Duals of a Family of $\mathcal{N} = 1$ Fixed Points*, *JHEP* **0608** (2006) 083, [[hep-th/0506206](#)].
- [19] O. Lunin and J. Maldacena, *Deforming field theories with $U(1) \times U(1)$ global symmetry and their gravity duals*, *JHEP* **0505** (2005) 033, [[hep-th/0502086](#)].
- [20] M. Kulaxizi, *On β -deformations and Noncommutativity*, [hep-th/0610310](#).
- [21] N. Seiberg and E. Witten, *String theory and noncommutative geometry*, *JHEP* **9909** (1999) 032, [[hep-th/9908142](#)].
- [22] V. Schomerus, *D -branes and Deformation Quantization*, *JHEP* **9906** (1999) 030, [[hep-th/9903205](#)].
- [23] J. Maldacena, *Probing near extremal black holes with D -branes*, *Phys. Rev. D* **57** (1998) 3736–3741, [[hep-th/9705053](#)].
- [24] J. Maldacena, *Branes Probing Black Holes*, *Nucl. Phys. Proc. Suppl.* **68** (1998) 17–27, [[hep-th/9709099](#)].
- [25] A. A. Tseytlin, *Born–Infeld action, supersymmetry and string theory*, [hep-th/9908105](#).
- [26] R. R. Metsaev and A. A. Tseytlin, *Supersymmetric $D3$ brane actions in $ads_5 \times S^5$* , *Phys. Lett. B* **436** (1998) 281–288, [[hep-th/9806095](#)].
- [27] I. Chepelev and A. A. Tseytlin, *Long–distance interactions of branes: correspondence between supergravity and super Yang–Mills descriptions*, *Nucl. Phys. B* **515** (1998) 73–113, [[hep-th/9709087](#)].

- [28] V. Balasubramanian, R. Gopakumar, and F. Larsen, *Gauge Theory, Geometry and the Large N Limit*, *Nucl. Phys. B* **526** (1998) 415–431, [[hep-th/9712077](#)].
- [29] V. V. Khoze, *Amplitudes in the beta-deformed Conformal Yang–Mills*, *JHEP* **0602** (2006) 040, [[hep-th/0512194](#)].
- [30] M. F. Sohnius, *Introducing supersymmetry*, *Phys. Rep.* **128** (1985) 39.
- [31] D. R. T. Jones, *Coupling constant reparameterization and finite field theories*, *Nucl. Phys. B* **277** (1986) 153.
- [32] A. Parkes and P. C. West, *Finiteness in rigid supersymmetric theories*, *Phys. Lett. B* **138** (1984) 99.
- [33] D. R. T. Jones and L. Mezincescu, *The chiral anomaly and a class of two-loop finite supersymmetric gauge theories*, *Phys. Lett. B* **138** (1984) 293.
- [34] A. J. Parkes and P. C. West, *Three-loop results in two-loop finite supersymmetric gauge theories*, *Nucl. Phys. B* **256** (1985) 340.
- [35] M. T. Grisaru, B. Milewski, and D. Zanon, *The structure of UV divergences in SS YM theories*, *Phys. Lett. B* **155** (1985) 357.
- [36] R. G. Leigh and M. J. Strassler, *Exactly marginal operators and duality in four dimensional $\mathcal{N} = 1$ supersymmetric gauge theory*, *Nucl. Phys. B* **447** (1995) 95, [[hep-th/9503121](#)].
- [37] V. A. Novikov, M. A. Shifman, A. I. Vainshtein, and V. I. Zakharov, *Beta Function in Supersymmetric Gauge Theories: Instantons Versus Traditional Approach*, *Phys. Lett. B* **166** (1986) 329.
- [38] M. A. Shifman and A. I. Vainshtein, *Solution of the Anomaly Puzzle in Susy Gauge Theories and the Wilson Operator Expansion*, *Nucl. Phys. B* **277** (1986) 456.
- [39] M. A. Shifman and A. I. Vainshtein, *On holomorphic dependence and infrared effects in supersymmetric gauge theories*, *Nucl. Phys. B* **359** (1991) 571.
- [40] O. Aharony and S. S. Razamat, *Exactly Marginal Deformations of $\mathcal{N} = 4$ SYM and of its Supersymmetric Orbifold Descendants*, *JHEP* **0205** (2002) 029, [[hep-th/0204045](#)].
- [41] S. S. Razamat, *Marginal Deformations of $\mathcal{N} = 4$ SYM and of its Supersymmetric Orbifold Descendants*, [hep-th/0204043](#). MSc thesis.
- [42] V. Niarchos and N. Prezas, *BMN operators for $N = 1$ superconformal Yang–Mills theories and associated string backgrounds*, *J. High Energy Phys.* **06** (2003) 015, [[hep-th/0212111](#)].
- [43] D. Z. Freedman and U. Gursoy, *Comments on the β -deformed $\mathcal{N} = 4$ SYM Theory*, *JHEP* **0511** (2005) 042, [[hep-th/0506128](#)].
- [44] A. Mauri, S. Penati, A. Santambrogio, and D. Zanon, *Exact results in planar $\mathcal{N} = 1$ superconformal Yang–Mills theory*, *JHEP* **0511** (2005) 024, [[hep-th/0507282](#)].
- [45] S. M. Kuzenko and A. Tseytlin, *Effective action of β -deformed $\mathcal{N} = 4$ SYM theory and AdS/CFT*, *Phys. Rev. D* **72** (2005) 075005, [[hep-th/0508098](#)].
- [46] G. C. Rossi, E. Sokatchev, and Y. S. Stanev, *New results in the deformed $\mathcal{N} = 4$ SYM theory*, [hep-th/0606284](#).
- [47] G. C. Rossi, E. Sokatchev, and Y. S. Stanev, *On the all-order perturbative finiteness of the deformed $\mathcal{N} = 4$ theory*, [hep-th/0606284](#).

- [48] S. Penati, A. Santambrogio, and D. Zanon, *Two point correlators in the beta-deformed $\mathcal{N} = 4$ SYM at the next-to-leading order*, *JHEP* **0510** (2005) 023, [[hep-th/0506150](#)].
- [49] A. Mauri, S. Penati, M. Pirrone, A. Santambrogio, and D. Zanon, *On the perturbative chiral ring for marginally deformed $\mathcal{N} = 4$ SYM theories*, *JHEP* **0608** (2006) 072, [[hep-th/0605145](#)].
- [50] F. Elmetti, A. Mauri, S. Penati, A. Santambrogio, and D. Zanon, *Conformal invariance of the planar beta-deformed $\mathcal{N} = 4$ SYM theory requires beta real*, [hep-th/0606125](#).
- [51] D. Berenstein and R. G. Leigh, *Discrete torsion, AdS/CFT and duality*, *J. High Energy Phys.* **0001** (2000) 038, [[hep-th/0001055](#)].
- [52] D. Berenstein, V. Jejjala, and R. G. Leigh, *Marginal and relevant deformations of $N = 4$ field theories and non-commutative moduli spaces of vacua*, *Nucl. Phys. B* **589** (2000) 196, [[hep-th/0005087](#)].
- [53] N. Dorey, T. J. Hollowood, and S. P. Kumar, *S-duality of the Leigh–Strassler deformation via matrix models*, *J. High Energy Phys.* **12** (2002) 003, [[hep-th/0210239](#)].
- [54] N. Dorey, *S-duality, deconstruction and confinement for a marginal deformation of $\mathcal{N} = 4$ SUSY Yang–Mills*, *J. High Energy Phys.* **0408** (2004) 043, [[hep-th/0310117](#)].
- [55] N. Dorey, *A new deconstruction of little string theory*, *J. High Energy Phys.* **0407** (2004) 016, [[hep-th/0406104](#)].
- [56] D. Berenstein and S. A. Cherkis, *Deformations of $N = 4$ SYM and integrable spin chain model*, *Nucl. Phys. B* **702** (2004) 49–85, [[hep-th/0405215](#)].
- [57] D. Bundzik and T. Mansson, *The general Leigh–Strassler deformation and integrability*, *JHEP* **0601** (2006) 116, [[hep-th/0512093](#)].
- [58] N. Beisert and R. Roiban, *Beauty and the Twist: The Bethe Ansatz for Twisted $\mathcal{N} = 4$ SYM*, *JHEP* **0508** (2005) 039, [[hep-th/0505187](#)].
- [59] S. Frolov, *Lax Pair for Strings in Lunin–Maldacena Background*, *JHEP* **0505** (2005) 069, [[hep-th/0503201](#)].
- [60] T. McLoughlin and I. Swanson, *Integrable twists in AdS/CFT*, *JHEP* **0608** (2006) 084, [[hep-th/0605018](#)].
- [61] L. F. Alday, G. Arutyunov, and S. Frolov, *Green–Schwarz Strings in TsT-transformed backgrounds*, *JHEP* **0606** (2006) 018, [[hep-th/0512253](#)].
- [62] O. Aharony, B. Kol, and S. Yankielowicz, *On exactly marginal deformations of $\mathcal{N} = 4$ SYM and Type IIB supergravity on $\text{AdS}_5 \times \text{S}^5$* , *JHEP* **0206** (2002) 039, [[hep-th/0205090](#)].
- [63] A. Catal-Ozer, *Lunin–Maldacena Deformations with Three parameters*, *JHEP* **0602** (2006) 026, [[hep-th/0512290](#)].
- [64] P. Sundell, *Spin($p+1, p+1$) Covariant Dp-brane Bound States*, *Int. J. Mod. Phys. A* **16** (2001) 3025–3040, [[hep-th/0011283](#)].
- [65] E. Bergshoeff, C. M. Hull, and T. Ortin, *Duality in the Type-II Superstring Effective Action*, *Nucl. Phys. B* **451** (1995) 547–578, [[hep-th/9504081](#)].
- [66] M. Cvetič, H. Lu, C. N. Pope, and K. S. Stelle, *T-duality in the Green–Schwarz Formalism and the Massless/Massive IIA Map*, *Nucl. Phys. B* **573** (2000) 149–176, [[hep-th/9907202](#)].
- [67] M. Fukuma, T. Oota, and H. Tanaka, *Comments on T-dualities of Ramond–Ramond Potentials*, *Prog. Theor. Phys.* **103** (2000) 425–446, [[hep-th/9907132](#)].

- [68] S. F. Hassan, *T-Duality, Space-time Spinors and R-R Fields in Curved Backgrounds*, *Nucl. Phys. B* **568** (2000) 145–161, [[hep-th/9907152](#)].
- [69] M. Kulaxizi and K. Zoubos, *Marginal Deformations of $\mathcal{N} = 4$ SYM from Open/Closed Twistor Strings*, *Nucl. Phys. B* **738** (2006) 317–349, [[hep-th/0410122](#)].
- [70] M. Kulaxizi and K. Zoubos, *Marginal Deformations of Tree-Level $\mathcal{N} = 4$ SYM from Twistor String Theory*, *Fortsch. Phys.* **54** (2006) 423–429, [[hep-th/0512302](#)].
- [71] S. A. Frolov, R. Roiban, and A. Tseytlin, *Gauge-string duality for superconformal deformations of $\mathcal{N} = 4$ Super Yang–Mills theory*, *JHEP* **0507** (2005) 045, [[hep-th/0503192](#)].
- [72] S. A. Frolov, R. Roiban, and A. Tseytlin, *Gauge-string duality for (non)supersymmetric deformations of $\mathcal{N} = 4$ Super Yang–Mills theory*, *Nucl. Phys. B* **731** (2005) 1–44, [[hep-th/0507021](#)].
- [73] H.-Y. Chen and S. P. Kumar, *Precision Test of AdS/CFT in Lunin–Maldacena Background*, *JHEP* **0603** (2006) 051, [[hep-th/0511164](#)].
- [74] C. Durnford, G. Georgiou, and V. Khoze, *Instanton test of non-supersymmetric deformations of the $AdS_5 \times S^5$* , *JHEP* **0609** (2006) 005, [[hep-th/0606111](#)].
- [75] G. Georgiou and V. Khoze, *Instanton Calculations in the beta-deformed AdS/CFT Correspondence*, *JHEP* **0604** (2006) 049, [[hep-th/0602141](#)].
- [76] R. Hernandez, K. Sfetsos, and D. Zoakos, *Gravity duals for the Coulomb branch of marginally deformed $\mathcal{N} = 4$ Yang–Mills*, *JHEP* **0603** (2006) 069, [[hep-th/0510132](#)].
- [77] U. Gursoy and C. Nunez, *Dipole deformations of $\mathcal{N} = 1$ SYM and Supergravity backgrounds with $U(1) \times U(1)$ global symmetry*, *Nucl. Phys. B* **725** (2005) 45–92, [[hep-th/0505100](#)].
- [78] C. Ahn and J. F. Vazquez-Poritz, *Marginal Deformations with $U(1)^3$ Global Symmetry*, *JHEP* **0507** (2005) 032, [[hep-th/0505168](#)].
- [79] C. Ahn and J. F. Vazquez-Poritz, *From Marginal Deformations to Confinement*, *JHEP* **0606** (2006) 061, [[hep-th/0603142](#)].
- [80] R. C. Rashkov, K. S. Viswanathan, and Y. Yang, *Generalizations of Lunin–Maldacena transformation on the $AdS_5 \times S^5$ background*, *Phys. Rev. D* **72** (2005) 106008, [[hep-th/0509058](#)].
- [81] D. Bundzik, *Star product and the general Leigh–Strassler deformation*, [hep-th/0608215](#).
- [82] J. Shelton, W. Taylor, and B. Wecht, *Nongeometric Flux Compactifications*, *JHEP* **0510** (2005) 085, [[hep-th/0508133](#)].
- [83] I. Ellwood and A. Hashimoto, *Effective descriptions of branes on non-geometric tori*, [hep-th/0607135](#).
- [84] D. Lowe, H. Nastase, and S. Ramgoolam, *Massive IIA string theory and matrix theory compactification*, *Nucl. Phys. B* **667** (2003) 55–89, [[hep-th/0303173](#)].
- [85] J. H. Schwarz, *Covariant Field Equations of Chiral $N=2$ $D=10$ Supergravity*, *Nucl. Phys. B* **226** (1983) 269.
- [86] C. G. Callan, C. Lovelace, C. R. Nappi, and S. A. Yost, *String Loop Corrections to Beta Functions*, *Nucl. Phys. B* **288** (1987) 525.
- [87] A. Abouelsaood, C. G. Callan, C. R. Nappi, and S. A. Yost, *Open Strings in Background Gauge Fields*, *Nucl. Phys. B* **280** (1987) 599.

- [88] A. S. Cattaneo and G. Felder, *A path integral approach to the Kontsevich quantization formula*, *Commun. Math. Phys.* **212** (2000) 591, [[math.QA/9902090](#)].
- [89] M. Kontsevich, *Deformation quantization of Poisson manifolds, I*, *Lett. Math. Phys.* **66** (2003) 157, [[q-alg/9709040](#)].
- [90] L. Cornalba and R. Schiappa, *Nonassociative star product deformations for D-brane worldvolumes in curved backgrounds*, *Comm. Math. Phys.* **225** (2002) 33, [[hep-th/0101219](#)].
- [91] M. Herbst, A. Kling, and M. Kreuzer, *Star Products from Open Strings in Curved Backgrounds*, *JHEP* **0109** (2001) 014, [[hep-th/0106159](#)].
- [92] M. Herbst, A. Kling, and M. Kreuzer, *Cyclicity of non-associative products on D-branes*, *JHEP* **0403** (2004) 003, [[hep-th/0312043](#)].
- [93] P.-M. Ho, *Making Non-Associative Algebra Associative*, *JHEP* **0111** (2001) 026, [[hep-th/0103024](#)].
- [94] R. Minasian, M. Petrini, and A. Zaffaroni, *Gravity duals to deformed SYM theories and Generalized Complex Geometry*, [hep-th/0606257](#).
- [95] I. Ellwood, *NS-NS fluxes in Hitchin's generalized geometry*, [hep-th/0612100](#).
- [96] J. de Boer, P. A. Grassi, and P. van Nieuwenhuizen, *Non-commutative superspace from string theory*, *Phys. Lett. B* **574** (2003) 98–104, [[hep-th/0302078](#)].
- [97] H. Ooguri and C. Vafa, *The C-Deformation of Gluino and Non-planar Diagrams*, *Adv. Theor. Math. Phys.* **7** (2003) 53–85, [[hep-th/0302109](#)].
- [98] N. Berkovits and N. Seiberg, *Superstrings in Graviphoton Background and $\mathcal{N} = 1/2 + 3/2$ Supersymmetry*, *JHEP* **0307** (2003) 010, [[hep-th/0306226](#)].